



Flint® Extra Fungicide

Group 11 Fungicide

Flint® Extra fungicide is a proven, dependable fungicide that offers enhanced value for growers.

/// BENEFITS

- **Proven Efficacy**
Provides proven broad-spectrum control of key diseases in various crops.
- **Excellent Compatibility**
Good crop safety history and mix compatibility with other products, making it an excellent fit for many Integrated Pest Management (IPM) programs.
- **Easy to Use**
Low use rates in an easy-to-use liquid 500 SC formulation.

/// THE POWER OF FLINT® EXTRA FUNGICIDE

The active ingredient in Flint Extra rapidly penetrates the waxy layer of the plant tissues and redistributes on the plant surface, providing locked-in protection against disease, even within hard-to-penetrate plant canopies.

The fungicide stops both early and developing fungal diseases by preventing initial infection as well as secondary spread of diseases. This makes Flint Extra a valuable tool in your disease management program.

/// FLINT® EXTRA FUNGICIDE QUICK FACTS


- **Active Ingredient:** Trifloxystrobin
- **FRAC:** Group 11 Fungicide
- **Single Word:** Caution
- **REI:** 12 Hours
- **Use Rate:** 2.0 to 4.7 fl. oz./A, depending on crop

/// RECOMMENDATION FOR USE IN GRAPES


In case of known Strobilurin resistance Bayer recommends a maximum of one solo Strobilurin application in a tank-mix. From a resistance management point of view, Strobilurins are still a valuable tool for growers since their use in a tank-mix and rotational program reduces the pressure on other fungicides (*FRAC Group 7, 3 etc.*).

Registered Crops	Key Diseases
<ul style="list-style-type: none">• Brassica Vegetables• Citrus• Cucurbit Vegetables• Fruiting Vegetables• Grapes• Hops• Leafy Vegetables• Pome Fruit• Potatoes• Rice• Root Vegetables• Stone Fruit• Strawberries• Sugar Beets• Tree Nuts	<ul style="list-style-type: none">• Alternaria• Anthracnose• Black Rot• Botryosphaeria• Botrytis• Cercospora Leaf Spot• Cherry Leaf Spot• Early Blight• Fly Speck• Greasy Spot• Melanose• Phomopsis• Plectosporium Blight• Post-Bloom Fruit Drop• Powdery Mildew• Rice Blast• Rust• Scab• Septoria• Shot Hole• Sooty Blotch• Stemphylium

In cherries, Flint® Extra Fungicide with Bravo Weather Stik® controlled powdery mildew and leaf spot.



In grapes, Luna Experience® Fungicide followed by Flint® Extra Fungicide controlled powdery mildew to trace levels.



Trial conducted near Walnut Grove by UCCE, University of California, Davis, 2016. Rates are on an oz. per acre basis. All treatments with adjuvant (not shown). Applications on 14-day interval. Bayer programs began with Microthiol® Disperss® 5 lb. up to bloom on 7-day interval. Other programs were alternated full season.

*Farmers living in Alaska, California or Hawaii do not qualify for Bayer PLUS Rewards

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

For more information, visit www.CropScience.Bayer.us.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Bayer, Bayer Cross, Flint® and Luna Experience® are registered trademarks of Bayer Group. All other trademarks are the property of their respective owners. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2023 Bayer Group. All rights reserved.

