



Luna

# Luna<sup>®</sup> Flex Fungicide Group 7 | 3 Fungicide - Apples

Luna<sup>®</sup> Flex fungicide is the newest member of the trusted Luna family of fungicides for the Eastern US. It offers long-lasting protection from key diseases in a variety of crops.

#### /// BENEFITS

#### • Broad Spectrum Labeled for key crops and diseases in the Eastern

US markets

Great Curative Control


Enhanced curative properties that will help improve overall disease control

Long-Lasting Protection
 Will provide extended performance over the recommended spray interval

### /// LUNA<sup>®</sup> FLEX FUNGICIDE QUICK FACTS


- Active Ingredient: Fluopyram and Difenoconazole
- FRAC: Group 7 and 3 fungicide
- Formulation Type: Liquid, 375 SC (Soluble Concentrate)
- Signal Word: Caution
- REI: 12 hours
- PHI: 14 days
- Use Rate: 8.0 oz./A

Key Apple Diseases
<ul> <li>Cedar Apple Rust</li> </ul>
<ul> <li>Flyspeck</li> </ul>
<ul> <li>Powdery Mildew</li> </ul>
• Scab
<ul> <li>Sooty Blotch</li> </ul>
<ul> <li>Various Rots</li> </ul>



# Luna® Flex Fungicide Effectively Controls Apple Scab on Foliage and Fruits

Apple McIntosh; Trial conducted in North Rose, NY in 2020 by an independent research firm. For head-to-head comparison fungicides were applied as a full season program with seven repeated foliar applications. Nonionic surfactant was used with all spray applications. Please follow the label for rates and timings for commercial applications. Data recorded at 24 days after last application for leaf and fruit scab, and 53 days after last application for fruit scab control in storage. Means within category followed by the same letter are not significantly different according to Fisher's Protected LSD (P<0.05)



# Luna<sup>®</sup> Flex Fungicide Effectively Controls Apple Scab

Results are the average of six trials conducted in 2021 in MI, NC, NY, OH, and PA. All treatments were three times in April-May; other fungicide applications in the programs were similar across all the treatments. Nonionic surfactant was used with all spray applications. Data was recorded in July-August. Means within category followed by the same letter are not significantly different according to Fisher's Protected LSD (P<0.05)

**IMPORTANT:** This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

Luna Flex contains oil in the formulation. If tank-mixing Luna Flex with oil-sensitive products, read and follow their label restrictions. Do not apply in tank-mix with, or 7 days before or after an application of Captan.

For more information, visit www.CropScience.Bayer.us.

Luna<sup>®</sup> FLEX

BAYER

#### ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registeration status in your state. Bayer, Bayer Cross, Luna Sensation®, Luna Tranquility® and Luna® Flex are registered trademarks of Bayer Group. All other trademarks are the property of their respective owners. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2023 Bayer Group. All rights reserved.