



**PLUS**

ELIGIBLE



**Luna<sup>®</sup>**  
**FLEX**

# Luna<sup>®</sup> Flex Fungicide

*Group 7 | 3 Fungicide - Strawberries*

Luna<sup>®</sup> Flex fungicide is the newest member of the trusted Luna family of fungicides for the Eastern US. It offers long-lasting protection from key diseases in a variety of crops.


### /// BENEFITS

- **Broad Spectrum**  
Labeled for key crops and diseases in the Eastern US markets
- **Great Curative Control**  
Enhanced curative properties to improve overall disease control
- **Long-Lasting Protection**  
Extended performance over the recommended spray interval


### /// LUNA<sup>®</sup> FLEX FUNGICIDE QUICK FACTS

- **Active Ingredient:** Fluopyram and Difenoconazole
- **FRAC:** Group 7 and 3 fungicide
- **Formulation Type:** Liquid, 375 SC (Soluble Concentrate)
- **Signal Word:** Caution
- **REI:** 12 hours
- **PHI:** 0 days
- **Use Rate:** 12.0 - 13.6 fl. oz./A

Other Key Labeled Crops	Key Strawberry Diseases
<ul style="list-style-type: none"> <li>• Apples</li> <li>• Bulb Vegetables</li> <li>• Bushberries</li> <li>• Citrus</li> <li>• Cucurbits</li> <li>• Fruiting Vegetables</li> <li>• Pome Fruits</li> <li>• Stone Fruits</li> </ul>	<ul style="list-style-type: none"> <li>• Anthracnose</li> <li>• Gray Mold</li> <li>• Leaf Rust</li> <li>• Leaf Spot</li> <li>• Powdery Mildew</li> <li>• Rhizopus Fruit Rot</li> </ul>



## Luna® Flex Fungicide Effectively Controls Strawberry Botrytis Fruit Rot



Trial conducted in Plant City, FL in 2021 by University of Florida. Test products were applied when weather conditions for Botrytis cinerea infection were favorable (17 to 25°C and ≥ 12 h leaf wetness) according to the Strawberry Advisory System. For head-to-head comparison test products were applied three times. Please follow the label for rates and timings for commercial applications. Other fungicide applications in the programs were similar across all the treatments. Yield was recorded based on the weight of marketable fruit and Botrytis fruit rot incidence was recorded as the percentage of symptomatic fruit relative to all harvested fruit. Means within series followed by the same letter are not significantly different according to Fisher's Protected LSD ( $P < 0.05$ ).



**IMPORTANT:** This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

**Luna Flex contains oil in the formulation. If tank-mixing Luna Flex with oil-sensitive products, read and follow their label restrictions. Do not apply in tank-mix with, or 7 days before or after an application of Captan.**

For more information, visit [www.CropScience.Bayer.us](http://www.CropScience.Bayer.us).

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Performance may vary, from location to location and from year to year, as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's fields. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Bayer, Bayer Cross, Luna Tranquility® and Luna® Flex are registered trademarks of Bayer Group. All other trademarks are the property of their respective owners. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at [www.BayerCropScience.us](http://www.BayerCropScience.us). Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2023 Bayer Group. All rights reserved.

**Luna®  
FLEX**

