

***FIFRA Section 24(c) Special Local Need Label*****FOR DISTRIBUTION AND USE ONLY WITHIN GEORGIA**

For Suppression of Alternaria Brown Spot and for control of Phytophthora Foot & Root Rot with a reduce Pre-Harvest Interval

ALIETTE WDG® Fungicide

EPA Reg. No. 264-516

SLN. No. GA-230002

THIS LABEL IS VALID UNTIL APRIL 30, 2024, UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

ACTIVE INGREDIENT: Aluminum tris (O-ethyl phosphonate)	80.0%
INERT INGREDIENTS:	20.0%
TOTAL:	100.0%

FIRST AID

IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Immediately call a poison control center or doctor for treatment advice.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Have person sip a glass of water if able to swallow.• Do not give anything by mouth to an unconscious person.
<p>For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.</p> <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****WARNING**

Harmful if swallowed or absorbed through the skin or inhaled. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic and estuarine invertebrates. Drift and runoff may be hazardous to aquatic and estuarine organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves (made of any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber), shoes plus socks, and protective eyewear.

CITRUS (BEARING)

ALIETTE WDG applied in conjunction with good cultural practices will provide effective control of foot rot, root rot, gummosis, and fruit brown rot caused by *Phytophthora* spp. ALIETTE WDG also controls ice-nucleating bacteria (California only) and suppresses Alternaria Brown Spot.

DISEASE	APPLICATION METHOD	RATE lbs product	APPLICATION PROGRAM
Phytophthora Foot Rot or Gummosis (<i>Phytophthora</i> spp.)	Trunk Spray or Paint	2.5 – 5.0/ 5 gals	Mix the desired amount of ALIETTE WDG with 5 gallons of water and apply to the trunk lesion in a sufficient volume to thoroughly wet the entire lesion. If no lesion is present, the application should be made from the soil line up the trunk approximately 2 feet. If trunk lesions are present the higher rate should be used.
Phytophthora Foot & Root Rot Brown Rot (<i>Phytophthora</i> spp.)	Foliar	5.0/Acre	Apply as a spray to wet when conditions favor disease development. (If you are unsure about whether these conditions exist, check with your Cooperative Extension Service.)
		1.0/100 gals	Apply as instructed above. This volumetric concentration is derived from a per acre foliar application rate of 5 lbs of ALIETTE WDG diluted in 500 gals of water.
Phytophthora Foot & Root Rot (<i>Phytophthora</i> spp.)	Chemigation	5.0/Acre	Begin application when conditions favor disease development. (If you are unsure about whether these conditions exist, check with your Cooperative Extension Service.) Inject ALIETTE WDG continuously during a scheduled irrigation.
Alternaria Brown Spot (<i>Alternaria</i> spp.) (Suppression only)	Foliar	5.0/Acre	ALIETTE WDG suppresses <i>Alternaria</i> spp. that cause Brown Spot. The first application should be applied in the spring at the first sign of new growth. A second and third application should follow at 30-45 day intervals. Apply in sufficient water and direct spray to insure thorough coverage of the susceptible foliage and fruit.
		1.0/100 gals	Apply as instructed above. This volumetric concentration is derived from a per acre foliar application rate of 5 lbs of ALIETTE WDG diluted in 500 gals of water.

RESTRICTIONS AND LIMITATIONS:

- Do not apply within 12 hours of harvest (PHI = 12 hours).
- Any combination of labeled application methods can be used, however, do not exceed a total of four applications or 20 lbs/Acre per year.
- For foliar applications do not exceed 500 GPA.
- ALIETTE WDG may be applied by ground, air, or chemigation.
- Do not graze livestock in treated groves.
- Do not feed forage from treated groves.

Note: In order to apply the correct amount of product to your groves you must know the number of gallons of water needed to spray one acre of trees to the point of drip. If you do not know this gallonage you should conduct a test to determine it. If you do not know how to conduct such a test with your equipment you should seek assistance from your equipment dealer or Cooperative Extension Service.

Note: Bayer CropScience has determined that phytotoxicity from solubilized copper may occur if products containing copper are tank-mixed with ALIETTE WDG or if unbuffered ALIETTE WDG is applied to foliage with copper residues.

Do not tank-mix with copper compounds. When applied prior to or after copper compounds, the pH of ALIETTE WDG should be raised to 6.0 or above with the addition of an alkaline buffer such as Potassium Carbonate (3 lbs Potassium Carbonate to 5 lbs ALIETTE WDG) or DiAmmonium Phosphate (5 lbs DiAmmonium Phosphate to 5 lbs ALIETTE WDG).

Adjuvants which enhance pesticide penetration may cause phytotoxicity when mixed with ALIETTE WDG.

Mixing ALIETTE WDG with surfactants or foliar fertilizers is not recommended. Citrus sensitivity to a particular combination should be determined by spraying a small amount of foliage. Evaluate 3-7 days later for adverse effects.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

**Bayer CropScience, LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167**