

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

GROUP	3	INSECTICIDE
-------	---	-------------

Baythroid® XL
EPA Reg. No. 264-840

For control of Brown Marmorated Stink Bug (BMSB) in Citrus, Grape and Potato.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ARIZONA AND CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

CITRUS

Crops of Crop Group 10 Including: Calamondin, Citrus citron, Citrus hybrids (Citrus spp., includes chironja, tangelo and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sweet and sour), Pummelo, Satsuma mandarin, White sapote (Casimiroa spp.), and other cultivars and/or hybrids of these.

Pest Controlled	Rate fluid oz/Acre	Rate lb AI/Acre
Brown Marmorated Stink Bug	3.2 – 6.4	0.025 – 0.05

Notes and Restrictions

Pre-Harvest Interval (PHI): **0 day.**

Maximum BAYTHROID XL allowed per 7-day interval: **6.4 fluid oz/A (0.05 lb AI/Acre).**

Maximum BAYTHROID XL allowed per crop season: **6.4 fluid oz/A (0.05 lb AI/Acre).**

Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.

Minimum application volume (water): **25 GPA – ground, 25 GPA – aerial application.**

GRAPE

Includes: Table grape, Raisin, Wine and Muscadine grape

PEST CONTROLLED	Rate fluid ounces/Acre	Rate lb AI/Acre
Brown Marmorated Stink Bug	2.4 – 3.2	0.019 – 0.025

Notes and Restrictions

Pre-Harvest Interval (PHI): **3 days.**

Maximum BAYTHROID XL allowed per 14-day interval: **3.2 fluid oz/A (0.025 lb AI/Acre).**

Maximum BAYTHROID XL allowed per crop season: **12.8 fluid oz/A (0.1 lb AI/Acre).**

Minimum application volume (water): 50 GPA – ground, 25 GPA – aerial application.

POTATO AND OTHER TUBEROUS AND CORM VEGETABLES

Includes all members of Crop Subgroup 1C:

Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Edible canna, Cassava (bitter and sweet), Chayote root, Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet potato, Tanier, True yam, Turmeric, Yam bean

PEST CONTROLLED	Rate fluid ounces/Acre	Rate lb AI/Acre
Brown Marmorated Stink Bug	1.6 – 2.8	0.013 – 0.022

Notes and Restrictions

Pre-Harvest Interval (PHI): **0 day.**

If more than 5.6 fluid oz/Acre is applied, allow at least 14 days between last application and grazing.

Maximum BAYTHROID XL allowed per 5-day interval: **2.8 fluid oz/A (0.022 lb AI/Acre).**

Maximum BAYTHROID XL allowed per crop season: **16.8 fluid oz/A (0.132 lb AI/Acre).**

For MEDICAL And TRANSPORTATION Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937)*