



Flexible Application Timing Helps Ensure Clean Fields at End-of-Season

Capreno® herbicide defends against tough weeds with excellent residual control, ensuring an amazingly clean finish.

/// BENEFITS

- **Broad-spectrum Weed Control**
Capreno® controls tough grass and broadleaf weeds, including those resistant to glyphosate, PPO, ALS, auxin and triazine chemistries.
- **Wide Application Window**
Capreno® has a wide application window and provides consistent performance across a variety of weather conditions.
- **Multiple Sites of Action**
Two effective sites of action and long-lasting residual activity help prevent the development of weed resistance.

/// FACTS

- Bayer's proprietary safener found in Capreno® enables plants to better withstand herbicidal activity.
- Capreno® can help manage resistance through the use of multiple herbicide sites of action if tankmixed with chemistries such as glyphosate and dicamba.
- Capreno® controls many weeds that exhibit resistance to glyphosate, ALS, PPO, auxin and triazine chemistries.

Registered Crops

- | | |
|---------------|--------------|
| • Field Corn | • White Corn |
| • Silage Corn | • Seed Corn |

Key Pests

- | | |
|------------------------|--------------------|
| • Foxtail Species | • Barnyardgrass |
| • Ragweed Species | • Palmer Amaranth |
| • Kochia | • Fall Panicum |
| • Common Lambsquarters | • Velvetleaf |
| • Common Cocklebur | • Common Waterhemp |

Formulation Information

- | | |
|------------------------------------|--------------------------|
| • Tembotrione (2.88 lbs ai/gal) | • Suspension Concentrate |
| • Thien carbazon (0.57 lbs ai/gal) | |

Application Rate	3 fl oz/A
Application Timing	Preemergence, Postemergence, (V1-20" or V7)
Site-of-Action Group	2 27
Recommended Tankmix Options	DiFlexx® herbicide 4 Atrazine 5 Degree® Xtra herbicide 5 15 Warrant® herbicide 15 Glyphosate 9

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

For more information, visit www.CropScience.Bayer.us.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Performance may vary, from location to location and from year to year, as local growing, soil and environmental conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on their growing environment. The recommendations in this material are based upon trial observations and feedback received from a limited number of growers and growing environments. These recommendations should be considered as one reference point and should not be substituted for the professional opinion of agronomists, entomologists or other relevant experts evaluating specific conditions. Degree Xtra® is a restricted use pesticide. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your state department(s) of agriculture, local dealer or representative for the product registration status in your state. Not all products are registered for use in all states and may be subject to use restrictions. Bayer, Bayer Cross, Capreno®, Degree Xtra®, DiFlexx®, and Warrant® Herbicide are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LLC, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2026 Bayer Group. All rights reserved.

Capreno

