Convintro™

Herbicide

Net Contents:

2.5 Gallons

DIFLUFENICAN | GROUP 12 | HERBICIDE

For preemergence control of certain broadleaf weeds on field corn and soybeans

ACTIVE INGREDIENT:	
Diflufenican [N-(2,4-difluorophenyl)-2-[3-(trifluoromethyl)phenoxy]pyridine-3-carboxamide]	**42.01%**
OTHER INGREDIENTS:	**57.99%**
TOTAL:	**100.00%**

Contains 4.17 pounds per U.S. gallon (500 grams per liter) of the active ingredient per U.S. gallon

EPA Reg. No. 264-1213

KEEP OUT OF REACH OF CHILDREN

CAUTION PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.

Please refer to booklet for additional precautionary statements and directions for use.

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours a Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

MIX WELL BEFORE USE

Produced for
Bayer CropScience LLC
800 N. Lindbergh Blvd.
St. Louis, MO 63167
Convintro is a trademark of Bayer Group.
©2026 Bayer Group. All rights reserved.

US91244017A 260630Av2 07/26

FIRST AID

If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. • **DO NOT** give anything by mouth to an unconscious person.
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

In case of emergency, call the toll-free Bayer CropScience Emergency Response telephone number: 1-800-334-7577.
Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

Note to Physician: No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- Harmful if swallowed or absorbed through the skin.
- Causes moderate eye irritation.
- Avoid contact with skin, eyes or clothing.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607 (d)(e) and (f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users must remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users must remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. For terrestrial uses, **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high

water mark. **DO NOT** apply when weather conditions favor drift from areas treated. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. **DO NOT** use the same spray equipment for other purposes unless thoroughly cleaned. **DO NOT** contaminate water used for irrigation or domestic purposes.

SURFACE WATER ADVISORY This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several months or more after application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ENDANGERED AND THREATENED SPECIES PROTECTION REQUIREMENTS:

Before using this product, you must obtain any applicable Endangered Species Protection Bulletins ('Bulletins') within six months prior to or on the day of application. To obtain Bulletins, go to Bulletins Live! Two (BLT) at https://www.epa.gov/pesticides/bulletins. When using this product, you must follow all directions and restrictions contained in any applicable Bulletin(s) for the area you are applying the product, including any restrictions on application timing if applicable. It is a violation of Federal law to use this product in a manner inconsistent with its labeling, including this labeling instruction to follow all directions and restrictions contained in any applicable Bulletin(s). For general questions or technical help, call 1-844-447-3813, or email ESPP@epa.gov.

For Important crop safety information, refer to the Use Directions section under the specific crop.

REPORTING ECOLOGICAL INCIDENTS: For guidance on reporting ecological incidents, see EPA's Pesticide Incident Reporting website: https://www.epa.gov/pesticideincidents.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval **(REI) of 12 hours.**

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

PRODUCT INFORMATION

Convintro Herbicide:

- is a selective herbicide for control of Amaranthus weed species infesting soybean, field corn, seed corn and corn grown for silage.
- is formulated as a suspension concentrate (SC) containing 4.17 pounds diflufenican active ingredient per gallon.
- contains the active ingredient diflufenican which is a PDS inhibitor mode of action that controls weeds by inhibiting enzymes that are essential to the protection of chlorophyll in plant leaves.
- is effective in controlling glyphosate-, triazine-, PPO-, ALS-, HPPD-, and auxin- herbicide resistant populations of Amaranthus weed species.

Requires the addition of tank mix partner(s) for control of weeds other than Amaranthus weed species.

USE RESTRICTIONS

DO NOT apply Aerially.

DO NOT use this product in the states of California, Alaska, or Hawaii.

DO NOT use this product in the state of Florida, except for the counties of: Escambia, Santa Rosa, Okaloos, Walton, Holmes, Washington, Jackson, Calhoun, Gadsden, Leon, Jefferson, Madison, Hamilton, Suwannee, Columbia, Baker, Lafayette, Union, Bradford, Clay, Gilchrist, Dixie, Levy, Alachua, Putnam, and Marion.

Refer to the specific use directions and restrictions in each specific crop section.

APPLICATION INSTRUCTIONS

Convintro Herbicide:

- may be used in either conventional, conservation tillage, or no-till crop management systems.
- may be applied either preplant or preemergence.
- will provide it's most effective weed control when applied and subsequently moved into the soil by rainfall, sprinkler irrigation or mechanical tillage prior to weed emergence.
- may be tankmixed or applied in sequential applications with other herbicides to control additional weeds.
- may be applied using either water carrier or sprayable grade fluid fertilizer as a liquid carrier.
- may be applied by ground application only. Aerial application is not permitted.
- may be applied as either a broadcast spray or as a band application.

Refer to the **SPECIFIC USE DIRECTIONS** section of this label for additional application information specific from each registered use of Convintro Herbicide.

Ground Application (Banding)

Banding herbicide application equipment must be carefully calibrated to prevent crop exposure to concentrations of Convintro Herbicide that exceed the labeled rate for the soil type. It is critical to ensure that the calibrated band width equates to actual band width realized in field applications. Bands actually delivered at a width narrower than targeted will concentrate the product and increase the risk for crop response.

Even flat spray tip nozzles and a band width of no less than 12" must be used.

Apply a broadcast equivalent rate and volume per acre. The following equations may be used to make the required calculations as follows:

$$\frac{\text{band width (inches)}}{\text{row width (inches)}} * \text{broadcast rate per acre} = \text{banding rate per acre}$$

$$\frac{\text{band width (inches)}}{\text{row width (inches)}} * \text{broadcast spray volume per acre} = \text{banding spray volume per acre}$$

Ground Application (Broadcast)

Apply Convintro Herbicide in tank mixtures in a minimum of 10 gallons of spray mixture per acre. Uniform, thorough spray coverage is important to achieve consistent weed control. Confirm that spray tank is equipped with a sparge tube agitator, mechanical agitation devices or jet agitator devices that are capable of providing vigorous agitation while mixing. Agitate thoroughly before and during application with either bypass or mechanical agitation. Maintain adequate in-tank agitation at all times, including during momentary stops to keep the spray mixture in suspension and provide a minimum of 20% bypass at all times. Keep the spray boom at the lowest possible spray height above the target surface. Refer to nozzle manufacturer's specifications for proper nozzle, pressure setting, and sprayer speed for optimum product performance and minimal spray drift. Uneven application or sprayers not properly calibrated may decrease the level of weed control and/or increase the level of adverse crop response. Maintain constant ground speed while applying product to ensure proper distribution. **DO NOT** overlap spray patterns beyond equipment manufacturers specifications as excessive rates may result in adverse crop responses and potential stand loss. The use of screens to protect the pump and nozzles is advised. Screens placed on the suction side of the pump must be 16-mesh or coarser. Equip sprayer with 50 mesh in line filter(s) located between the pump discharge and the spray boom. This filter may be located prior to each spray boom section. If nozzle filters are used, use 50 mesh or coarser nozzle filters as specified by nozzle manufacturer. **DO NOT** place a screen in the recirculation line. Application must be made at a sufficient spray pressure and volume to provide accurate and uniform application of spray particles to a given area without causing spray drift to non-target areas.

RESISTANCE MANAGEMENT

For resistance management, Convintro Herbicide is a Group 12 herbicide. Any weed population may contain or develop plants naturally resistant to Convintro Herbicide and other Group 12 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of Convintro Herbicide or other Group 12 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- **DO NOT** apply this product alone. An appropriate herbicide tank mix partner must be used to broaden the spectrum of weed control.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

MANDATORY SPRAY DRIFT MITIGATIONS

MANDATORY SPRAY DRIFT MANAGEMENT

- **DO NOT apply this product aerially.**

Ground Boom Applications

- During application, the Sustained Wind Speed, as defined by the National Weather Service (standard averaging period of 2 minutes), must register between 3 and 15 miles per hour.
- Wind speed and direction must be measured on location using a windsock or anemometer (including systems to measure wind speed or velocity using application equipment).
- Wind speed must be measured at the release height or higher, in an area free from obstructions such as trees, buildings, and farm equipment.
- **DO NOT** release spray at a height greater than 2 feet above the ground or crop canopy.
- Applicators must select nozzle and pressure that deliver coarse or coarser depending on team discussions] droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASAE S572).
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. Be aware of nearby non-target sites and environmental conditions.

- IMPORTANCE OF DROPLET SIZE
  An effective way to reduce spray drift is to apply large droplets. Consider the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.
- Controlling Droplet Size - Ground Boom
  - Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Consider using the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
  - Pressure – Using the lowest spray pressure recommended for the nozzle will produce the target spray volume and droplet size.
  - Spray Nozzle - Consider using a spray nozzle that is designed for the intended application as well as using nozzles designed to reduce drift.
- Release height – Ground Boom
  For ground equipment, the boom should remain level with the crop and have minimal bounce. Automated boom height controllers are recommended with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.
- HOODED (OR SHIELDED) SPRAYERS
  Shielding the boom or individual nozzles can reduce spray drift. Consider using hooded sprayers. Applicators should verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY
  When making applications in hot and dry conditions, consider using larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS
  Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.
- WIND
  Drift potential generally increases with wind speed.
  Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

MANDATORY RUNOFF/EROSION MITIGATION

Applicators must access and search Bulletins Live! Two (BLT) at https://www.epa.gov/pesticides/bulletins within six months prior to or on the day of the application to determine whether the application site falls within a Pesticide Use Limitation Area (PULA) that has a Bulletin in BLT. If you are located inside a PULA, follow the instructions in the bulletin.

If the application site is located outside a PULA, follow the instructions in the section below.

2 mitigation points are required for all field crops listed on this label. Follow the steps below to achieve points.

- Step 1. To achieve the mitigation points specified above, visit EPA's mitigation menu website (www.epa.gov/pesticides/mitigation-menu) for a full list of mitigation and mitigation relief options.
- Step 2. Determine if you are eligible for mitigation relief. Runoff/erosion mitigation is NOT required if certain field/application parameters are present at the time of application (*e.g.*, subsurface or tile drains with controlled outlet, perimeter berm systems, irrigation tailwater return systems, spot treatment, etc).
- Step 3. If the application site does not meet the field/application parameters specified on EPA's mitigation menu website, choose among the mitigation and/or mitigation relief options on EPA's mitigation menu website to meet or exceed the required points noted here before applying this product.
- Step 4. To achieve mitigation points for the application, the mitigation and mitigation relief measures must be:

- Employed in accordance with the instructions and descriptions on EPA's Mitigation Menu Website.
- In place during the application unless a different timing (such as before or after application) is specifically provided in the measure's description on EPA's Mitigation Menu Website. EPA may periodically update the Mitigation Menu Website, for example, by adding new mitigation measures or updating a mitigation measure description.

- Step 5. Additional restrictions may be present in bulletins—always follow the most restrictive bulletin instructions. If you are located in an area where PULAs overlap, follow the most restrictive requirements across all bulletins. When tank mixing, the most restrictive requirements must be followed between the products' labels and bulletins.

COMPATIBILITY TESTING AND TANK MIX PARTNERS

Compatibility

If Convintro Herbicide is to be tank mixed with liquid fertilizers or other pesticides, compatibility must be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 qt) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, **DO NOT** use this mixture for spraying. Indications of incompatibility usually will appear within 5-15 minutes after mixing. Read and follow all parts of the label of each tank-mix product.

Order of Mixing

Convintro Herbicide may be used with other recommended pesticides, fertilizers, and micronutrients. When using Convintro Herbicide, make sure the sprayer is completely clean, free of rust or corrosion which occurs from winter storage. Examine strainers and screens to be sure the sprayer is clean from previously used pesticides. Ensure that all in-line strainer and nozzle screens in the sprayer are 50 mesh or coarser. **DO NOT** use screens finer than 50 mesh.

Any tank mix containing Convintro Herbicide must be kept agitated and sprayed out immediately. **DO NOT** allow tank mixes to stand for prolonged periods of time.

The proper mixing procedure for Convintro Herbicide, in tank mix combinations with other pesticides is:

1. Fill the spray tank with 50% of the required volume of water and begin agitation. Agitation must continue throughout the entire mixing and spraying procedure.
2. If a compatibility agent is necessary to improve mixing or to prevent formation of undesirable and unsprayable gels or precipitates, while agitating, add it to the carrier already in the tank. Use only compatibility agents cleared by FDA for this use. Read and follow directions for use, cautionary statements, and all other information appearing on the selected compatibility agent label. Check for adequate agitation.
3. Add specified rate of Convintro Herbicide slowly to the spray tank or mixing system and ensure complete dispersion. Maintain and ensure thorough dispersion and sufficient agitation during both mixing and spraying.
4. Add the tank mix product.
5. Fill the spray tank with the balance of water needed.
6. Continue agitation during application and until sprayer tank is empty.

RE-SUSPENDING SC PRODUCTS IN SPRAY SOLUTION

Like other suspension concentrates (SCs), Convintro Herbicide will settle if left standing without agitation. If the spray solution is allowed to settle for one hour or more, reagitate the spray solution for a minimum of 10 minutes before application.

APPLICATION OF CONVINTRO HERBICIDE IN FLUID FERTILIZERS

Convintro Herbicide may be applied in fluid fertilizer solutions by following the appropriate mixing procedures and compatibility check. When using tank mix combinations, be sure all components are compatible.

Compatibility checks of Convintro Herbicide and tank mix combinations which include Convintro Herbicide must be made for each batch of fluid fertilizer because of the variability of these fertilizers.

Compatibility Check:

1. Pre-mix 2 teaspoonfuls of Convintro Herbicide with 16 teaspoonfuls of water (1:8 ratio) in a quart jar by adding the water first and follow with Convintro Herbicide. Mix thoroughly. Add the second herbicide after mixing Convintro Herbicide first.
2. Then pour 1 pint of fluid fertilizer into the quart jar and shake well.
3. Allow to stand for 5 minutes.

THIS COMPATIBILITY CHECK MUST ONLY BE USED WHEN MIXING WITH FLUID FERTILIZERS.

Interpretation of Results: If the solution in the jar appears to be uniform, without signs of agglomeration, or without a separation of an oily film on top of the fertilizer, the mixture may be used. If not, repeat the compatibility check using twice the amount of water or add a compatibility agent to the water. If separation occurs, but the mixture can be resuspended by shaking, then application is possible with good agitation in the spray tank.

Tank mixing Guidelines:

1. Add the required amount of water and compatibility agent (if required) to the tank. Start the agitation system while adding Convintro Herbicide followed by a second herbicide and agitate.
2. Add fluid fertilizer and continue agitation.
3. Maintain continuous agitation to assure uniform spray mixture until the tank is emptied.

EQUIPMENT CLEANOUT PROCEDURES

To avoid injury or exposure to non-target crops, thoroughly clean all mixing and spray equipment, including pumps, nozzles, lines and screens with a good quality tank cleaner, on an approved rinse pad or on the field site where an approved crop is to be grown. Mix only as much cleaning solution as needed.

1. Flush tank, hoses, boom, and nozzles with clean water.
2. Use a pressure washer with a high quality commercial spray tank cleaner in water to clean the inside of the spray tank. Take care to wash all parts of the tank, including the inside top surface. If a pressure washer is not available, completely fill the sprayer with the cleaning solution to ensure contact of the cleaning solution with all internal surfaces of the tank and plumbing. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.
3. Flush hoses, spray lines, and nozzles for at least 1 minute with the cleaning solution.
4. Dispose of rinsate from steps 1-3 in an appropriate manner.
5. Repeat steps 2-4.
6. Remove nozzles, screens, and strainers and clean separately in the cleaning solution after completing the above procedures.
7. Rinse the complete spraying system with clean water.

Cleanup must be conducted on an approved rinse pad or the field site where an approved crop is to be grown.

ROTATIONAL CROPS

Rotational crops vary in their crop response to low concentrations of Convintro Herbicide remaining in the soil. The amount of Convintro Herbicide that may be present in the soil depends on soil moisture, soil temp, application rate, elapsed time since application, and other environmental factors. When Convintro Herbicide is used in combination with other products, always follow the most restrictive rotational crop requirements. The following rotational crops may be planted after applying Convintro Herbicide: Labeled crops Corn (field and seed) and Soybeans may be replanted at any time; all other crops require a minimum plant back interval of 30 days, except legumes at a minimum of 365 days.

In the event of crop failure: If the crop treated with Convintro Herbicide is lost, only soybeans, field corn, and corn grown for silage may be replanted immediately. **DO NOT** make an additional application of Convintro Herbicide.

Crop	Geography	Rotational Interval (months)	Minimum Precipitation Requirement[1]
Corn (Field and seed) & Soybeans	All	0	None
Wheat, Triticale, Barley, and Rye	All	4	
Sweet corn, Popcorn, Potato, Grain Sorghum, and Sunflower	All	6	
Cotton and Rice	All	10	15 inches of cumulative precipitation from application to planting of rotational crop.[2]
Sugar beets	East of the Mississippi River		
Tobacco, Alfalfa, Canola, Chickpeas, Peas, Peanuts and Lentils	All	12	
Dry Beans	East of the Mississippi River	12	
Dry Beans, Sugar beets	West of the Mississippi River	18	
All other crops[3]	All		

[1] The amount of cumulative precipitation required before planting a rotational crop is in addition to the required rotational interval given in months.

[2] Furrow or flood irrigation not to be included in total. No more than 7 inches of overhead irrigation included in total.

[3] All other crops may be seeded only after the completion of a successful bioassay after a Convintro Herbicide application. Refer to the "Field/Small Scale Bioassay" section.

Cover Crops

Use of cover crops as a means of soil improvement, erosion control, weed and/or insect suppression, etc., following harvest of corn or soybeans in the Fall is increasing. Planting of cover crops in fields treated with Convintro Herbicide is allowed as long as these cover crops are not grazed by livestock nor harvested for food. Cover crops are to be tilled under or chemically controlled with burndown herbicides in the spring. Many cover crops can be planted within 90-120 days after application of Convintro Herbicide.

However, all potential cover crops have not been evaluated for tolerance to Convintro Herbicide and significant injury may occur. Prior to seeding a cover crop, complete a successful field/ home bioassay to provide an indication of the level of tolerance to the prior Convintro Herbicide application. Refer to the "Field/Small Scale Bioassay" section. If used in tank mixtures with other herbicides, always follow the most restrictive label.

Field/Small Scale Bioassay

A field/ small scale bioassay must be completed before rotating to a cover crop other than those specified in the "Rotational Crop Restrictions" section of this label. To conduct an effective **field bioassay**, grow strips of the crop(s) you intend to grow the following season in a field previously treated with Convintro Herbicide. The test strip must be placed in a controlled area and include low areas and knolls, and include variations in soil including type and pH. Crop response to the bioassay will determine if the crop(s) grown in the test strips can be grown safely in the areas previously treated with Convintro Herbicide.

For an effective **small scale bioassay**, collect uniform samples of all soil types from the Convintro Herbicide- treated field (see example above for types of soil in the sample) and place the soil into a sturdy container. Plant the desired cover crop into the soil, apply water and place the container in a warm, sunny area to allow germination and growth of the crop. Monitor growth of the cover crop over a three to four week period. If the cover crop emerges and grows normally, the risk to establish and grow the cover crop in the Convintro Herbicide-treated field must be tolerable.

WEEDS CONTROLLED

Convintro Herbicide applied as directed in this label will control or suppress the weeds listed below. Additional weeds may be controlled with tank mixtures or sequential applications (refer to the TANK MIX INSTRUCTIONS section of this label). Always refer to the tank mix partner labels for specific use rates and additional instructions.

BROADLEAF WEEDS CONTROLLED				
Broadleaf Weeds	Convintro Herbicide Alone		Broadleaf Weeds	Convintro Herbicide Alone
Amaranth, Palmer	C		Waterhemp, common	C
Pigweed, prostrate	C		Waterhemp, tall	C
Pigweed, redroot	C			
Pigweed, smooth	C			
C= Control				

SPECIFIC USE DIRECTIONS

CORN (Field Corn, Seed Corn and Corn Grown for Silage)

Convintro Herbicide may be used in either conventional, conservation tillage, or no-till crop management systems and may be applied either preplant or preemergence. Convintro Herbicide treatments are most effective in controlling weeds when adequate rainfall is received within 14 days after application. If cultivation is necessary because of soil crusting, soil compaction, or weed germination before rain occurs, use shallow tillage including rotary hoe to lightly incorporate Convintro Herbicide. Make certain corn seeds are below the tilled area. If treated soil is moved during tillage practices in such a way that the herbicide barrier is no longer intact, weeds may emerge from areas where treated soil has been removed. **DO NOT** incorporate with a drag harrow after planting. An appropriate herbicide tank mix partner must be used in order to broaden spectrum of weed control.

APPLICATION TIMING

Preplant Surface-Applied

Convintro Herbicide may be applied up to 30 days before planting corn when used in a planned sequential application program including Convintro Herbicide followed by postemergence applied herbicides appropriate for control of the target weeds. An appropriate herbicide tank mix partner must be used to broaden spectrum of weed control. Total Convintro Herbicide applied may not exceed the listed rate for a preplant treatment on the predominate soil type in the field.

Moving treated soil out of the row or moving untreated soil to the surface during planting may result in reduced weed control. For enhanced burndown of emerged broadleaf and grass weeds tank mix with glyphosate, dicamba, paraquat, or 2,4-D according to their labelled use rates. Follow all label directions, rotational crop guidelines, and use restrictions/precautions for Convintro Herbicide and any tank-mix partners and use the most restrictive guidelines.

Preemergence

Apply Convintro Herbicide during planting (behind the planter after furrow closure) or within 3 days after planting. Failure to thoroughly close and firm the seed furrow may allow herbicide to directly contact the seed which can cause injury. An appropriate tank mix partner herbicide must be used to broaden spectrum of control. For enhanced burndown of emerged broadleaf and grass weeds tank mix with glyphosate, dicamba, paraquat, or 2,4-D according to their labelled use rates. If applying Convintro Herbicide to emerged weeds, the inclusion of an appropriate adjuvant (Crop Oil Concentrate (COC), Methylated Seed Oil (MSO), or High Surfactant Oil Concentrate (HSOC)) will enhance burndown activity of controlled weeds. Follow all label directions, rotational crop guidelines, and use restrictions/precautions for Convintro Herbicide and any tank-mix partners and use the most restrictive guidelines.

Application in fall or early spring following harvest of previous crop

Convintro Herbicide may be applied after previous crop harvest and at least 30 days prior to planting corn using the rates in the preplant section. For enhanced burndown of emerged broadleaf and grass weeds tank mix with glyphosate, dicamba, paraquat, or 2,4-D according to their labelled use rates. If applying SC500 Herbicide to emerged weeds, the inclusion of an appropriate adjuvant (Crop Oil Concentrate (COC), Methylated Seed Oil (MSO), or High Surfactant Oil Concentrate (HSOC)) will enhance burndown activity of controlled weeds. Follow all label directions, rotational crop guidelines, and use restrictions/precautions for Convintro Herbicide and any tank-mix partners and use the most restrictive guidelines.

APPLICATION RATES

Application Timing	Maximum Rate in Fluid Ounces of Convintro Herbicide per Acre					
	Coarse Soils		Medium Soils		Fine Soils	
	Sand, Loamy Sand, Sandy loam		Loam, Silt Loam, Silt, Sandy Clay Loam		Silty Clay Loam, Clay Loam, Sandy Clay, Silty Clay, Clay	
	< 1.5% O.M.*	>=1.5% O.M.	< 1.5% O.M.	>=1.5% O.M.	< 1.5% O.M.	>=1.5% O.M.
Preplant 8 to 30 days prior to planting	2.4 (0.0784 lbs ai/A)	3.3 (0.1076 lbs ai/A)			4.1 (0.1334 lbs ai/A)	
Preemerge (0 to 7 days prior to planting or within 3 days of planting)	2.4 (0.0784 lbs ai/A)		3.3 (0.1076 lbs ai/A)		4.1 (0.1334 lbs ai/A)	

*Organic Matter

Application of Convintro Herbicide at less than listed rates for the appropriate soil will only provide suppression of sensitive weeds.

RESTRICTIONS FOR USE

- **DO NOT** apply more than 4.1 fluid ounces per acre (0.1334 lbs) of Convintro Herbicide per 365-day period or exceed the maximum labeled rate for any given soil type.
- **DO NOT** apply more than 4.1 fluid ounces (0.1334 lbs) per acre of Convintro Herbicide in a single application or exceed the maximum labeled rate for any given soil type or crop response may occur.
- **DO NOT** make more than one application per year.
- **DO NOT** apply under conditions that favor runoff
- **DO NOT** apply if soil is saturated with water or above field capacity.

PRECAUTIONS FOR USE

- Spray overlaps produce areas of over application which increase the potential for crop injury.
- **Planting depth:** Seed must be planted a minimum of 1 ½ inches deep and must be completely covered with soil and furrow firmed or reduced crop stand or injury may occur.
- **Application Timing:** Applying Convintro Herbicide less than 8 days before corn planting may increase the risk of crop injury, especially if irrigation or rainfall is received following the application. Injury symptoms include leaf tissue bleaching (whitening), leaf chlorosis (yellowing), necrosis (browning), and/or temporary stunting. Applying Convintro Herbicide more than 8 days (but not more than 30 days) prior to planting may reduce the risk of crop injury.
- **Effect of adverse weather:** Following an application of Convintro Herbicide, periods of cool/cold, wet conditions (cool/cold daytime/nighttime temperatures, saturated soil conditions, recurring rainfall events, etc.) during corn seed germination and/or early crop development period may result in temporary crop injury. Injury symptoms may appear as leaf tissue bleaching (whitening) and/or crop stunting. Corn plants usually recover from this injury without affecting yield.
- **Effect of variable soils on use rate:** The proper use rate of Convintro Herbicide is affected by several soil factors, including soil texture, organic matter, and soil pH. Soils which contain variations in one or more of these factors in a given area are termed variable soils and may be more likely to incur localized corn injury symptoms from an application of Convintro Herbicide. Characteristics of localized soil variants that are more likely to incur injury are a coarser soil texture, a lower organic matter and/or a higher pH (alkaline/calcareous soil) than other areas of the same field and include, among others, clay knolls, eroded hill sides, and terracing with scraped exposed subsoil. The user is responsible for selecting the rate of Convintro Herbicide that is appropriate for all soils in the area of application.

- **Corn hybrids and certain male pollinators:** Corn hybrids and certain male pollinators within blended corn varieties vary in their response to Convintro Herbicide. Not all hybrids or male pollinators within blended corn varieties have been tested for sensitivity to Convintro Herbicide. You should consult with your seed provider, your local Bayer CropScience representative, and/or other knowledgeable agricultural professionals for advice on hybrids or varieties containing male pollinator lines before applying Convintro Herbicide. If the tolerance of a hybrid or variety containing a male pollinator line is not known, you must apply Convintro Herbicide to a small area to first determine if the hybrid is tolerant prior to spraying large acreages of that hybrid.

SPECIAL PRECAUTIONS

Injury to corn may occur when Convintro Herbicide is used under the following conditions:

- When soils have a calcareous surface area or a pH of 7.5 or higher.
- When applied in conjunction with soil-applied organophosphate pesticides.
- Over application or boom overlapping may result in stand loss and soil residues.
- Uneven application can decrease the level of weed control and/or increase the level of injury.
- When applied to any soil with less than ½% organic matter.
- When sprayers are not calibrated accurately.
- When heavy rains occur soon after application, especially in poorly drained areas where water may stand for several days.
- When corn is planted less than 1.5 inches deep, particularly in preemergence application.

Activation: A minimum amount of soil moisture is required to activate Convintro Herbicide. In areas of low rainfall, preemergence applications to dry soil must be followed with light irrigation of 1/4 acre inch of water. **DO NOT** apply heavy irrigation immediately after application. As with many surface-applied herbicides, weed control and crop tolerance may vary with rainfall and/or soil texture.

Replanting: If replanting is necessary in fields treated with Convintro Herbicide as directed on this label, the field may be replanted to corn or soybeans (depending on the tank mix herbicide utilized). When replanting a minimum of tillage is recommended. **DO NOT** apply a second treatment as injury to corn may occur.

TANK MIX INSTRUCTIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Convintro Herbicide must be used in tank mixture with other herbicides for improved control of certain broadleaf and grass weeds in corn. Preplant/preemergence tank mixtures with Convintro Herbicide may include but are not limited to those listed. Refer to and follow all parts of the label of each tank-mix partner. Tank-mix combinations may be used in either conventional, conservation tillage, or no-till cropping systems and applied at the same timings as Convintro Herbicide unless otherwise specified in the tank mix label. Multiple tank mixtures are allowed unless otherwise specified by the respective product labels. Check all tank-mix product labels for proper rates and compatibilities for multiple tank-mixes.

DO NOT TANK MIX ANY PRODUCT CONTAINING ISOXAFLUTOLE WITH THIS PRODUCT.

Possible Preplant/Preemergence Tank Mix Partners for Additional Weed Control

DiFlexx® (EPA Reg. No. 264-1173; *dicamba*)
Roundup PowerMAX® 3 (EPA Reg. No. 524-659; *glyphosate*)
RT 3® (EPA Reg. No. 524-544; *glyphosate*)
Harness® (EPA Reg. No. 524-473; *acetochlor*)
Degree Xtra® (EPA Reg. No. 524-511; *acetochlor*, *atrazine*)
Harness® Xtra (EPA Reg. No. 524-480; *acetochlor*, *atrazine*)
Harness® Xtra 5.6L (EPA Reg. No. 524-485; *acetochlor*, *atrazine*)
Harness® Max (EPA Reg. No. 524-636; *acetochlor*, *mesotrione*)

2,4-D	dicamba	glufosinate-ammonium	paraquat
flumetsulam	fluthiacet-methyl	glyphosate	clopyralid
dimethenamid-P	acetochlor	s-metolachlor	pyroxasulfone
	flufenacet	thiencarbazone-methyl	atrazine

SOYBEAN

Convintro Herbicide may be used in either conventional, conservation tillage, or no-till crop management systems and may be applied either preplant or preemergence. Convintro Herbicide treatments are most effective in controlling weeds when adequate rainfall is received within 14 days after application. If cultivation is necessary because of soil crusting, soil compaction, or weed germination before rain occurs, use shallow tillage including rotary hoe to lightly incorporate Convintro Herbicide. Make certain soybean seeds are below the tilled area. If treated soil is moved during tillage practices in such a way that the herbicide barrier is no longer intact, weeds may emerge from areas where treated soil has been removed. **DO NOT** incorporate with a drag harrow after planting. An appropriate tank mix partner herbicide must be used in order to broaden spectrum of weed control.

APPLICATION TIMING

Preplant Surface-Applied

Convintro Herbicide may be applied up to 30 days before planting soybean when used in a planned sequential application program including Convintro Herbicide followed by postemergence applied herbicides appropriate for control of the target weeds. An appropriate herbicide tank mix partner must be used in order to broaden spectrum of weed control. Total Convintro Herbicide applied may not exceed the listed rate for a preplant treatment on the predominant soil type in the field. Moving treated soil out of the row or moving untreated soil to the surface during planting may result in reduced weed control. For enhanced burndown of emerged broadleaf and grass weeds tank mix with glyphosate, dicamba, paraquat, or 2,4-D according to their labelled use rates. Follow all label directions, rotational crop guidelines, and use restrictions/precautions for Convintro Herbicide and any tank-mix partners and use the most restrictive guidelines.

Preemergence

Apply Convintro Herbicide during planting (behind the planter after furrow closure) or within 3 days after planting. Failure to thoroughly close and firm the seed furrow may allow herbicide to directly contact the seed which can cause injury. An appropriate tank mix herbicide partner must be used in order to broaden spectrum of weed control. For enhanced burndown of emerged broadleaf and grass weeds tank mix with glyphosate, dicamba, paraquat, or 2,4-D according to their labelled use rates. If applying Convintro Herbicide to emerged weeds, the inclusion of an appropriate adjuvant (Crop Oil Concentrate (COC), Methylated Seed Oil (MSO), or High Surfactant Oil Concentrate (HSOC)) will enhance burndown activity of controlled weeds. Follow all label directions, rotational crop guidelines, and use restrictions/precautions for Convintro Herbicide and any tank-mix partners and use the most restrictive guidelines.

Application in fall or early spring following harvest of previous crop

Convintro Herbicide may be applied after previous crop harvest and at least 30 days prior to planting soybean using the rates in the preplant section. For enhanced burndown of emerged broadleaf and grass weeds tank mix with glyphosate, dicamba, paraquat, or 2,4-D according to their labelled use rates. If applying SC500 Herbicide to emerged weeds, the inclusion of an appropriate adjuvant (Crop Oil Concentrate (COC), Methylated Seed Oil (MSO), or High Surfactant Oil Concentrate (HSOC)) will enhance burndown activity of controlled weeds. Follow all label directions, rotational crop guidelines, and use restrictions/precautions for Convintro Herbicide and any tank-mix partners and use the most restrictive guidelines.

<table>
<tr><th rowspan="3">Application Timing</th><th colspan="6">Maximum Rate in Fluid Ounces of Convintro Herbicide per Acre</th></tr>
<tr><th colspan="2">Coarse Soils

Sand, Loamy Sand, Sandy Loam</th><th colspan="2">Medium Soils

Loam, Silt Loam, Silt, Sandy Clay Loam</th><th colspan="2">Fine Soils

Silty Clay Loam, Clay Loam, Sandy Clay, Silty Clay, Clay</th></tr>
<tr><th>< 1.5% O.M.*</th><th>>=1.5% O.M.</th><th>< 1.5% O.M.</th><th>>=1.5% O.M.</th><th>< 1.5% O.M.</th><th>>=1.5% O.M.</th></tr>
<tr><td>Preplant 8 to 30 days prior to planting</td><td>2.4 (0.0784 lbs ai/A)</td><td colspan="2">3.3 (0.1076 lbs ai/A)</td><td colspan="2">4.1 (0.1334 lbs ai/A)</td><td>4.9 (0.1597 lbs ai/A)</td></tr>
<tr><td>Preemerge (0 to 7 days prior to planting or within 3 days of planting)</td><td colspan="2">2.4 (0.0784 lbs ai/A)</td><td>3.3 (0.1076 lbs ai/A)</td><td colspan="2">4.1 (0.1334 lbs ai/A)</td><td>4.1 - 4.9 (0.1334 lbs - 0.1597 lbs ai/A)</td></tr>
</table>

*Organic Matter

Application of Convintro Herbicide at less than listed rates for the appropriate soil will only provide suppression of sensitive weeds.

RESTRICTIONS FOR USE

- **DO NOT** apply more than 4.9 fluid ounces per acre (0.1597 lbs) of Convintro Herbicide per 365-day period or exceed the maximum labeled rate for any given soil type.
- **DO NOT** apply more than 4.9 fluid ounces (0.1597 lbs) per acre of Convintro Herbicide in a single application or exceed the maximum labeled rate for any given soil type or crop response may occur.
- **DO NOT** make more than one application per year.
- **DO NOT** apply under conditions that favor runoff
- **DO NOT** apply if soil is saturated with water or above field capacity.

PRECAUTIONS FOR USE

- Spray overlaps produce areas of over application which increase the potential for crop injury.
- **Planting depth:** Soybean seed must be planted a minimum of 1 ½ inches deep and must be completely covered with soil and furrow firmed or reduced crop stand or injury may occur.
- **Application Timing:** Applying Convintro Herbicide less than 8 days before soybean planting may increase the risk of crop injury, especially if irrigation or rainfall is received following the application. Injury symptoms may appear as leaf tissue bleaching (whitening), leaf chlorosis (yellowing), necrosis (browning), and/or temporary crop stunting. Applying Convintro Herbicide more than 8 days (but not more than 30 days) prior to planting may reduce the risk of crop injury.
- **Effect of variable soils on use rate:** The proper use rate of Convintro Herbicide is affected by several soil factors, including soil texture, organic matter, and soil pH. Soils which contain variations in one or more of these factors in a given area are termed variable soils and may be more likely to incur localized soybean injury symptoms from an application of Convintro Herbicide. Characteristics of localized soil variants that are more likely to incur injury are a more coarse soil texture, a lower organic matter and/or a higher pH (alkaline/calcareous soil) than other areas of the same field and include, among others, clay knolls, eroded hill sides, and terracing with scraped exposed subsoil. The user is responsible for selecting the rate of Convintro Herbicide that is appropriate for all soils in the area of application.
- **Effect of adverse weather:** Following an application of Convintro Herbicide, extended periods of cool/cold, wet conditions (cool/cold daytime/nighttime temperatures, saturated soil conditions, recurring rainfall events, etc.) during soybean seed germination and/or early crop development period may result in temporary crop injury. Injury symptoms may appear as leaf tissue bleaching (whitening) and/or crop stunting. Soybean plants usually recover from this injury without affecting yield.

SPECIAL PRECAUTIONS

Injury to soybeans may occur when Convintro Herbicide is used under the following conditions:

- When soils have a calcareous surface area or a pH of 7.5 or higher.
- When applied in conjunction with soil-applied organophosphate pesticides.
- Over application or boom overlapping may result in stand loss and soil residues.
- Uneven application can decrease the level of weed control and/or increase the level of injury.
- When applied to any soil with less than ½% organic matter.
- When sprayers are not calibrated accurately.
- When heavy rains occur soon after application, especially in poorly drained areas where water may stand for several days.
- When soybeans are planted less than 1 ½ inches deep, particularly in preemergence application.

Activation: A minimum amount of soil moisture is required to activate Convintro Herbicide. In areas of low rainfall, preemergence applications to dry soil must be followed with light irrigation of 1/4 acre inch of water. **DO NOT** apply heavy irrigation immediately after application. As with many surface-applied herbicides, weed control and crop tolerance may vary with rainfall and/or soil texture.

Rate Ranges: Where a rate range is shown, use a lower rate on soils that are coarse-textured or low in organic matter. Use a higher rate on soils that are relatively fine-textured or high in organic matter.

Replanting: If replanting is necessary in fields treated with Convintro Herbicide as directed on this label, the field may be replanted to corn (if 4.1 fluid ounces per acre or less was applied) or soybeans. Minimize tillage when replanting. **DO NOT** apply a second treatment as injury to soybeans may occur.

TANK MIX INSTRUCTIONS

TANK MIX INSTRUCTIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Convintro Herbicide may be used in tank mixture with other herbicides for improved control of certain broadleaf and grass weeds in soybeans. Preplant/preemergence tank mixtures with Convintro Herbicide include but are not limited to those listed. Refer to and follow all parts of the label of each tank-mix partner. Tank-mix combinations may be used in either conventional, conservation tillage, or no-till cropping systems and applied at the same timings as Convintro Herbicide unless otherwise specified in the tank mix label. Multiple tank mixtures are allowed unless otherwise specified by the respective product labels. Check all tank-mix product labels for proper rates and compatibilities for multiple tank-mixes.

DO NOT TANK MIX ANY PRODUCT CONTAINING ISOXAFLUTOLE WITH THIS PRODUCT.

Possible Preplant/Preemergence Tank Mix Partners for Additional Weed Control

Roundup PowerMAX® 3 (EPA Reg. No. 524-659; *glyphosate*)			
RT 3® (EPA Reg. No. 524-544; *glyphosate*)			
Warrant® (EPA Reg. No. 524-591; *acetochlor*)			
acetochlor	dicamba	glufosinate-ammonium	paraquat
glyphosate	pyroxasulfone	s-metolachlor	dimethenamid-P
pendimethalin	trifluralin	flufenacet	sulfentrazone
saflufenacil	flumioxazin	2,4-D	chlorimuron-ethyl
cloransulam-methyl	metribuzin		

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. **DO NOT** walk through spilled material. Dispose of pesticide as directed above. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response Telephone No. is 1-800-334-7577.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site in accordance with the uses permitted on this label, or at an approved waste disposal facility.

Container Disposal

Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons)

Non-refillable plastic container. **DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Once container is rinsed, offer for recycling if available, or if allowed by State or local authorities, puncture and dispose of in a sanitary landfill.

IMPORTANT NOTICE – PLEASE READ: LIMITATIONS OF WARRANTIES, LIABILITY, AND REMEDIES

This Notice of Limitation of Warranties, Liability, and Remedies ("Notice") and instructions to the purchaser and/or user ("Purchaser") contained in this product ("Product") label, including without limitation under Directions for Use (collectively, "Directions for Use"), are included in the terms of sale of this Product. Please read the Directions for Use and this Notice entirely before using this Product. The Purchaser accepts, acknowledges, and agrees to be bound by the Directions for Use and the terms of this Notice upon use of the Product. If Purchaser does not accept such terms, Purchaser must return the unopened Product container immediately. Any use and/or transfer of this Product must be authorized by Bayer CropScience LLC and accompanied by this Notice.

INHERENT RISKS OF USE: The Directions for Use of this Product are believed to be adequate, and Purchaser must carefully follow the Directions for Use. However, it is impossible to eliminate all risks associated with the use of this Product. Crop injury, ineffectiveness, or other unintended consequences may result because of factors and conditions beyond the control of Bayer CropScience LLC and its authorized Product distributors ("Seller"), including, among other things, adverse weather conditions, presence of other materials, and the manner of use or application. To the extent consistent with applicable law, Purchaser assumes all such risks.

To the extent the Product is a seed treatment product, Purchaser acknowledges that treatment of damaged seed (including, without limitation, highly mechanically damaged seed) or seed of low vigor or poor quality may result in reduced germination or seed and seedling vigor. Prior to use of this Product, Purchaser should inspect seed for damage and treat and conduct germination tests on a small portion of seed before treating a full seed lot with any seed treatment product.

EXPRESS WARRANTY: Seller's sole and exclusive warranty ("Exclusive Warranty") on the Product is the statements made on this Product label.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, EXCEPT FOR THE EXCLUSIVE WARRANTY SET FORTH ABOVE, SELLER DISCLAIMS ANY AND ALL WARRANTIES WITH RESPECT TO THIS PRODUCT, WHETHER EXPRESS OR IMPLIED (EITHER IN FACT OR BY OPERATION OF LAW), INCLUDING BUT NOT LIMITED TO: (A) THE IMPLIED WARRANTY OF MERCHANTABILITY; (B) THE IMPLIED WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE; (C) THE IMPLIED WARRANTY AGAINST NONINFRINGEMENT (FOR THIS PRODUCT ALONE OR IN COMBINATION WITH ANY OTHER PRODUCTS); AND (D) ANY WARRANTIES OF CROP PERFORMANCE OR, IF APPLICABLE, CARRYOVER SEED PERFORMANCE.

LIMITATION OF LIABILITY AND REMEDIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW:

1. SELLER'S TOTAL LIABILITY AND PURCHASER'S EXCLUSIVE REMEDY FOR ANY AND ALL LOSSES, INJURIES AND/OR DAMAGES ARISING FROM THE PURCHASE, USE, OR HANDLING OF THIS PRODUCT, OR OTHERWISE ARISING OUT OF A BREACH BY SELLER OF THE EXCLUSIVE WARRANTY, HOWEVER SUCH LIABILITY MAY ARISE, WHETHER SUCH CLAIMS ARE BASED ON CONTRACT, NEGLIGENCE, STRICT LIABILITY, TORT, OR ANY OTHER THEORY OF RECOVERY OR REMEDY, SHALL BE, AT THE ELECTION OF SELLER OR SELLER'S DELEGATE, AN AMOUNT NOT TO EXCEED THE PURCHASE PRICE PAID BY PURCHASER FOR THIS PRODUCT (AS SET FORTH IN THE APPLICABLE INVOICE) OR THE REPLACEMENT OF THE PRODUCT.
2. SELLER SHALL NOT BE LIABLE TO PURCHASER AND/OR ANY THIRD PARTY FOR ANY INCIDENTAL, CONSEQUENTIAL, RELIANCE, REMOTE, EXEMPLARY, PUNITIVE, SPECIAL, OR INDIRECT DAMAGES INCURRED OR EXPENDED IN THE PURCHASE, USE OR HANDLING OF THIS PRODUCT.
3. PURCHASER AGREES THAT IF THE PURCHASE PRICE PAID BY PURCHASER FOR THIS PRODUCT OR REPLACEMENT PRODUCT IS PROVIDED, THE REMEDY SET FORTH IN THIS NOTICE WILL NOT HAVE FAILED OF ITS ESSENTIAL PURPOSE.

PROMPT NOTICE OF CLAIMS REQUIRED: To the extent consistent with applicable law, as a condition to receiving Purchaser's limited remedy set forth above, any and all claims brought against the Seller must be brought within 30 days after the condition or event giving rise to the claim is discovered or should have been discovered, or prior to the harvest of any crop to which the Product was applied, whichever comes first, so that the claim can be investigated, and the Product or crop inspected.

MISCELLANEOUS: Purchaser agrees that this Notice is the entire agreement between Seller and Purchaser regarding Seller's warranty and liability for this Product. No modification of, addition to, or waiver of any of the terms of this Notice shall be binding unless set forth in writing and signed by an authorized representative of Bayer CropScience LLC. If any portion of this Notice not material to the remaining portions shall be held illegal, void, or ineffective by a governmental authority, the remaining portions shall remain in full force and effect. If any portion of this Notice is in conflict with any applicable statute or rule of law, then such portion shall be deemed to be modified to conform to such statute or rule of law.

Bayer (reg'd), Bayer Cross (reg'd), Convintro™ Herbicide, Roundup PowerMAX® 3, Warrant®, DiFlexx®, Harness®, Balance Flexx®, Corvus®, TriVolt®, Degree Xtra®, Harness® Xtra, Harness® Xtra 5.6L, Harness® Max, and Xtendimax® are trademarks of Bayer Group.
©2026 Bayer Group. All rights reserved.

Bayer

Convintro™ Herbicide

DIFLUFENICAN | GROUP 12 | HERBICIDE

For preemergence control of certain broadleaf weeds on field corn and soybeans

ACTIVE INGREDIENT:	
Diflufenican [N-(2,4-difluorophenyl)-2-[3-(trifluoromethyl)phenoxy]pyridine-3-carboxamide]	**42.01%**
OTHER INGREDIENTS:	**57.99%**
TOTAL:	**100.00%**

Contains 4.17 pounds per U.S. gallon (500 grams per liter) of the active ingredient per U.S. gallon

EPA Reg. No. 264-1213

KEEP OUT OF REACH OF CHILDREN

CAUTION PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.

Please refer to booklet for additional precautionary statements and directions for use.

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours a Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

MIX WELL BEFORE USE

FIRST AID

If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. • **DO NOT** give anything by mouth to an unconscious person.
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

In case of emergency, call the toll-free Bayer CropScience Emergency Response telephone number: 1-800-334-7577.
Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

Note to Physician: No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

- Harmful if swallowed or absorbed through the skin.
- Causes moderate eye irritation.
- Avoid contact with skin, eyes or clothing.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove and wash contaminated clothing before reuse.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. **DO NOT** walk through spilled material. Dispose of pesticide as directed above. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response Telephone No. is 1-800-334-7577.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site in accordance with the uses permitted on this label, or at an approved waste disposal facility.

Container Disposal

Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons)

Non-refillable plastic container. **DO NOT** reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Once container is rinsed, offer for recycling if available, or if allowed by State or local authorities, puncture and dispose of in a sanitary landfill.

Bayer (reg'd) and Convintro™ Herbicide are trademarks of Bayer Group. ©2026 Bayer Group. All rights reserved.

Produced for
Bayer CropScience LLC
800 N. Lindbergh Blvd.
St. Louis, MO 63167

Bayer

US91244017A 260630Av2 07/26

NET CONTENTS: 2 1/2 GALLONS

1 97515 27521 1