



RESTRICTED USE PESTICIDE

May injure (phytotoxic) susceptible non-target plants.

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Commercial and certified applicators must ensure that all persons involved in these activities are informed of the precautionary statements.

ISOXAFLUTOLE	GROUP	27	HERBICIDE
THIENCARBAZONE-METHYL	GROUP	2	HERBICIDE

FIFRA Section 24(c) Special Local Need (SLN) Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEW MEXICO

For Fall or Spring Applications to Control Kochia, Puncturevine and Russian Thistle in Fallow or Eco-Fallow Fields In UNION, QUAY, CURRY, ROOSEVELT and LEA Counties Only.

CORVUS[®] Herbicide

EPA REG No. 264-1066

EPA SLN No. NM-150002

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the EPA registered label.

Product	Rate	Crop	Weeds
CORVUS Herbicide	3-4 fl. oz/acre	Fallow, Eco-Fallow	For enhanced control of kochia, puncturevine, and Russian thistle

- Apply CORVUS Herbicide in the fall or spring to stubble of harvested crops or fallow/eco-fallow fields for control.
- **DO NOT** apply to frozen soils.
- Apply as a broadcast spray with ground equipment using a minimum of 10 gallons of spray mixture per acre.
- CORVUS Herbicide should be tank mixed with additional herbicides labeled for use in fallow or eco-fallow applications in order to achieve improved residual weed control. Recommended tank mix partners include atrazine (0.5-1.0 lb. ai/acre) or metribuzin (0.19-0.375 lb. ai/acre).
- For enhanced burndown of emerged broadleaf or grasses weeds, tank mix the CORVUS Herbicide + atrazine or metribuzin mixture with a labelled burndown herbicide such as dicamba, 2,-4-D, Gramoxone[®] or glyphosate (Note: glyphosate will not add additional control of glyphosate-resistant weeds). Always include an appropriate adjuvant (COC, MSO, HSOC) and nitrogen source (UAN, AMS) for enhanced burndown activity.
- Follow all label directions, rotational crop guidelines and use restrictions/precautions for CORVUS Herbicide and any tank-mix partners and use the most restrictive guidelines.

RESTRICTIONS / PRECAUTIONS

• USE IN COARSE TEXTURED SOILS WITH A SHALLOW WATER TABLE:

If the water table (i.e., level of saturation) is less than 25 feet below the ground surface, do not use on soils meeting all three of the following criteria (if less than three criteria are met or if the water table is greater than 25 feet below the ground surface, there is no restriction against application):

- The surface soil texture is sandy loam, loamy sand or sand.
- The subsoil texture is loamy sand or sand.
- The average organic matter (in the upper 12 inches) is less than 2% by weight.

ROTATIONAL CROP RESTRICTIONS

Rotational crops vary in their response to low concentrations of CORVUS Herbicide remaining in the soil. The amount of CORVUS Herbicide that may be present in the soil depends on soil moisture, soil temperature, application rate, elapsed time since application and other environmental factors. When CORVUS Herbicide is used in combination with other products, always follow the most restrictive rotational crop requirements. The following rotational crops may be planted after applying CORVUS Herbicide:

Minimum plant back intervals for various crops following CORVUS Herbicide.

Crop	Rotational Interval ²	Minimum Precipitation Requirement ¹
Field corn	0 Months	None
Wheat, Triticale, Cereal and Rye	4 Months	None
Barley, Soybean, Sweet corn ³ , Popcorn ³	9 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Rice ³ , Cotton ³	10 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Peanuts ³	11 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Tobacco ³	12 Months	15 inches of cumulative precipitation from application to planting of rotational crop
Alfalfa, Green and Dry Beans, Oats, Sorghum ⁴ , Sunflower, Canola, Potato, Sugar beet and all other crops ⁵	17 Months ³	30 inches of cumulative precipitation from application to planting of rotational crop

¹ The amount of cumulative precipitation required before planting a rotational crop is in addition to the required rotational interval given in months. Furrow or flood irrigation must not be included in total. No more than 7 inches of overhead irrigation must be included in total.

² Crop varieties planted back at intervals of one year or less must not have known acute sensitivity to ALS-inhibiting and/or SU herbicides.

³ When soil pH is 7.5 or above, crop plant back needs to be delayed to 17 months and to 24 months for crops listed in the 17 month interval above.

⁴ For CORVUS Herbicide used at 2.25 - 3.3 fl oz. per acre or less and the total of Thiencarbazone-methyl from all sources is 0.014 pounds active ingredient per acre or less, sorghum can be planted at the 9 month or longer interval.

⁵ All other crops may be seeded only after the completion of a successful bioassay after a CORVUS Herbicide application. Refer to the "Field/Small Scale Bioassay" section of the EPA registered label.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LLC
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)