



# Raising the Bar on Yields

Delaro® 325 SC fungicide for corn and soybeans can help you get the edge you're looking for. Delaro 325 SC has a broad spectrum of disease control and best-in-class, dual mode of action residual. Plus, it can improve plant health.

## /// BENEFITS

- **Broad-Spectrum Disease Control**  
Formulated with two heavy-hitting modes of action, together providing excellent control of even the toughest diseases.
- **Improved Plant Health**  
Delaro 325 SC promotes healthy, dark green leaves for increased photosynthesis and increases plant stress resistance to utilize the full genetic potential and yield of your seed.
- **Best-in-Class Dual Mode of Action Residual**  
Delaro 325 SC delivers long-lasting residual control from both its strobilurin and triazole components for extended performance from each active ingredient against important foliar diseases.

## /// FACTS

Delaro 325 SC maximizes and extends its protection of your crop by diffusing in and on the waxy surface of the plant. It then expands protection across the plant's exterior by moving via surface moisture. Within the plant, Delaro 325 SC is transported systemically throughout plant tissue for enhanced coverage.

Delaro 325 SC provides excellent preventative and curative defense against yield-robbing diseases which can improve harvestability.

### Registered Crops

- |   |  |
|---|--|
| <ul style="list-style-type: none"> <li>• Barley</li> <li>• Chickpeas</li> <li>• Corn, Field, for Seed</li> <li>• Corn, Sweet, for Seed</li> <li>• Cotton</li> </ul> | <ul style="list-style-type: none"> <li>• Dry Peas</li> <li>• Flax</li> <li>• Lentil</li> <li>• Soybean</li> <li>• Wheat</li> </ul> |
|---|--|

### Key Corn Diseases

- Anthracnose Leaf Blight
- Eyespot
- Gray Leaf Spot
- Northern corn Leaf Blight
- Northern corn Leaf Spot
- Physoderma Brown Spot
- Rust
- Southern Corn Leaf Blight
- Tar Spot

### Key Soybean Diseases

- Alternaria Leaf Spot
- Anthracnose
- Asian Soybean Rust
- Brown Spot
- Cercospora Blight
- Frogeye Leaf Spot
- Pod and Stem Blight
- Powdery Mildew
- Rhizoctonia Aerial Blight
- White Mold (suppression)

### Formulation Information

- Prothioconazole (1.49 lbs ai/gal)
- Trifloxystrobin (1.27 lbs ai/gal)



### /// APPLICATION TIMING

This versatile fungicide has application flexibility in corn — at-tassel timing (VT to R2), in combination with a herbicide (V4 to V8) and/or in between (V8 to VT).

In soybeans, the ideal time to first apply Delaro 325 SC is prior to development of most common diseases, between R1 and R3, with applications repeated as necessary. To manage white mold, apply prior to disease development, at R1, with a second treatment at R3.



### /// APPLICATION METHODS

Delaro 325 SC fungicide can be applied by air, ground or by chemigation and can be mixed with adjuvants, herbicides, insecticides and micronutrients for cost efficiency. Review the Delaro 325 SC fungicide label for complete application instructions.

### /// APPLY BEFORE DISEASE APPEARS

Consider planning ahead to protect your crop yield potential even if disease symptoms are not yet visible. Between V4 and V7, corn plants go through a critical growth period that greatly influences yield potential and increased stress tolerance.

<b>Application Rate</b>	Corn (Field Corn, Popcorn and Seed Corn) V4 to V7: 4 - 6 fl oz/A V8 to R2: 8 - 12 fl oz/A  Soybean R1 to R4: 8 - 11 fl oz/A
<b>Site-of-Action Group*</b>	<ul style="list-style-type: none"> <li>Prothioconazole <b>3</b></li> <li>Trifloxystrobin <b>11</b></li> </ul>
<b>Recommended Tankmix Options</b>	The inclusion of an adjuvant in the spray tank, for applications made through V8 (the collar of the eighth leaf is visible) and after tassel emergence (VT) is recommended.



\*For more information on fungicide group numbers and site-of-action options, visit [IWillTakeAction.com](http://IWillTakeAction.com)

**IMPORTANT:** This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

**DELARO®**



LEARN MORE

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.

Performance may vary, from location to location and from year to year, as local growing, soil and environmental conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on their growing environment. The recommendations in this material are based upon trial observations and feedback received from a limited number of growers and growing environments. These recommendations should be considered as one reference point and should not be substituted for the professional opinion of agronomists, entomologists or other relevant experts evaluating specific conditions. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Delaro® is a registered trademark of Bayer Group. ©2025 Bayer Group. All rights reserved.

