



PROTHIOCONAZOLE	GROUP	3	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE
FLUOPYRAM	GROUP	7	FUNGICIDE

FIFRA Section 24(c) Special Local Need Label
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

Delaro® Complete

For Cercospora Leaf Spot Control in Sugar Beets and White Mold in Dried Beans

EPA Reg. No. 264-1207

SLN. No. ID-260002

THIS LABEL IS VALID UNTIL AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

SUGAR BEET		
FOLIAR DISEASE CONTROLLED	RATE PER APPLICATION fl oz/acre	APPLICATION INSTRUCTIONS
Cercospora Leaf Spot (<i>Cercospora beticola</i>)	8.0 – 11.0 (0.092 - 0.126lb/acre prothioconazole) (0.081 - 0.111 lb/acre trifloxystrobin) (0.067 – 0.092 lb/acre fluopyram)	Apply DELARO COMPLETE Fungicide at the first sign of disease. Apply as needed on a 14 to 21 day spray schedule, as long as the maximum annual rate is not exceeded. Use the higher specified use rate and shorter specified intervals when conditions are favorable for severe disease pressure and/or when growing less disease resistant varieties.
Restrictions: - Maximum single application rate: [11.0 fl oz/acre of DELARO COMPLETE (0.126 lb/acre prothioconazole, 0.111 lb/acre trifloxystrobin and 0.092 lb/acre of fluopyram).] - Maximum annual application rate: [33.0 fl oz of DELARO COMPLETE (0.379 lb/acre prothioconazole, 0.333 lb/acre trifloxystrobin and 0.276 lb/acre fluopyram).] - Maximum number of applications: [3 (at 8.0 - 11.0 fl oz/acre).] - Pre-Harvest Interval (PHI): 21 day(s) - Minimum application volumes: 10 gallons/Acre (Ground); 5 gallons/Acre (Aerial) - Regardless of formulation or type of application, DO NOT apply more than 0.53 lbs prothioconazole or 0.33 lbs trifloxystrobin or 0.446 lbs fluopyram per acre per year from all uses, including seed treatment, soil and foliar applications.		
DRIED SHELLED PEA AND BEAN – EXCEPT SOYBEAN (Subgroup 6C), EXCEPT PEA		
Bean; Bean (<i>Lupinus</i> spp.) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); Bean (<i>Phaseolus</i> spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); Bean (<i>Vigna</i> spp.) (includes adzuki bean, blackeye pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); Broad bean (dry) (fava bean); Guar; Lablab bean (hyacinth bean).		
FOLIAR DISEASE CONTROLLED	RATE PER APPLICATION fl oz/acre	APPLICATION INSTRUCTIONS
White mold (<i>Sclerotinia sclerotiorum</i>)	11.7 (0.134 lbs prothioconazole/A + 0.118 lbs/acre Trifloxystrobin + 0.098 lbs/acre fluopyram)	Begin fungicide applications preventatively and continue as needed on a 10 to 14 day interval, as long as the maximum annual rate is not exceeded. Use the shorter specified intervals when conditions are favorable for severe disease pressure.
Restrictions: - Maximum single application rate: 11.7 fl oz/acre of DELARO COMPLETE (0.134 lb prothioconazole/acre + 0.118 lbs trifloxystrobin/acre + 0.098 lbs fluopyram/acre). - Maximum annual application rate: 23.4 fl oz/acre of DELARO COMPLETE (0.268 lb prothioconazole/acre + 0.235 lbs trifloxystrobin/acre + 0.195 lbs fluopyram/acre). - Maximum number of applications: 2 - Pre-Harvest Interval (PHI): 30 day(s) - DO NOT allow livestock to graze forage or hay and DO NOT harvest forage or hay for food or feed. - Minimum application volumes: 10 gallons/Acre (Ground); 5 gallons/Acre (Aerial) - Regardless of formulation or method of application, DO NOT apply more than 0.534 lbs prothioconazole, or 0.238 lbs trifloxystrobin, or 0.446 lbs fluopyram per acre per year from all uses, including seed treatment, soil and foliar applications.		

PROTHIOCONAZOLE	GROUP	3	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE
FLUOPYRAM	GROUP	7	FUNGICIDE

USE RESTRICTIONS:

- Do not use with handheld application equipment, including mechanically pressurized spray gun, backpack or tank pressurized spray gun or handheld boom applicators.
- Under certain conditions conducive to extended infection periods, additional fungicide applications beyond the number allowed by this label may be needed. Under these conditions, use another fungicide registered for the crop/disease.
- Do not use more than two consecutive applications of DELARO COMPLETE or any other Group 11 or Group 7 QoI fungicide without alternating with a different fungicide group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

**Bayer CropScience LLC
800 N. Lindbergh Blvd.
St. Louis, MO 63167**