



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

FLINT® Extra

Reduced rate in Apple for certain diseases
when used in tank-mix with protectant

EPA Reg. No. 264-826

FIFRA 2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF CONNECTICUT,
DELAWARE, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, MINNESOTA,
NEW HAMPSHIRE, NEW JERSEY, NEW YORK, PENNSYLVANIA, RHODE
ISLAND, VERMONT, VIRGINIA, WEST VIRGINIA, WISCONSIN**

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

APPLE		
Disease Controlled	Product Rate (fl oz/A)	Product Instructions
Scab (<i>Venturia</i> spp.)	2.0	Begin applications at green tip and continue on a 7- to 10-day interval as needed. Use with a protectant (e.g., Sulfur, Captan, Mancozeb, Serenade) tank mix depending on local disease control needs. Always use labeled rates for mix partners.
Cedar Apple Rust (<i>Gymnosporangium juniperivirginianae</i>)	2.0	Apply on a 7- to 10-day interval as needed. Alternate (every other application) with a sterol inhibitor fungicide. Use with a protectant (e.g., Sulfur, Captan, Mancozeb, Serenade) tank mix depending on local disease control needs. Always use labeled rates for mix partners.
Fly Speck (<i>Schizothyrium pomi</i>) Powdery mildew (<i>Podosphaera leucotricha</i>) Sooty Blotch (<i>Gloeodes pomigena</i>)	2.0	Apply on a 10- to 14-day interval as needed. Alternate (every other application) with a sterol inhibitor fungicide. Use with a protectant (e.g., Sulfur, Captan, Mancozeb, Serenade) tank mix depending on local disease control needs. Always use labeled rates for mix partners.
Application Restrictions: Pre-Harvest Interval (PHI): 14 day(s) Minimum interval between applications: 7 days Maximum FLINT Extra allowed per year: 10.5 fluid ounces/Acre - To reduce the potential for resistance, limit Group 11 fungicides to two sequential applications and alternate with at least two applications of fungicides from a different Group before making a third application with a Group 11 fungicide. - Do not apply more than 4 applications of FLINT Extra or any other Group 11 fungicide per year. - Do not apply FLINT Extra where spray drift may reach Concord grapes or crop injury may occur. Spray equipment must be rinsed after applying FLINT Extra before application of other products to Concord grapes or crop injury may occur.		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*