



Huskie®

Proven Broadleaf Control plus Resistance Management

Huskie® Herbicide is a powerful tool for resistance management that effectively controls a wide spectrum of broadleaf weeds, including bio-types resistant to herbicide Groups 4 and 9.

/// BENEFITS

- **Broad-Spectrum Control**
Controls a wide spectrum of tough-to-control broadleaf weeds.
- **Weed Resistance Management**
Includes herbicide Groups 6 and 27 to help fight weed resistance.
- **Tankmix Flexibility**
Excellent tankmix partner with grass herbicides to fit the needs of your farm.

/// APPLICATION TIMING AND RATES

- Apply Huskie herbicide to actively growing wheat, barley, rye or triticale between 1 leaf and up to flag leaf emergence. For grain sorghum, Huskie may be applied between 3 leaf stage of growth up to 30 inches and/or prior to flag leaf emergence, whichever comes first.
- Best results are obtained when applications are made to young actively growing weeds.

Registered Crops

- | | |
|----------------|----------------|
| • Spring Wheat | • Winter Wheat |
| • Barley | • Sorghum |
| • Durum | |

Key Weeds Controlled

- | | |
|-------------------|--------------------------|
| • Kochia | • Russian Thistle |
| • Palmer Amaranth | • Pigweed |
| • Prickly Lettuce | • Ragweed |
| • Buckwheat | • Nightshade** |
| • Lambsquarters | • Other Broadleaf Weeds* |

* See back for full list

**Controlled in spring wheat, winter wheat, durum and Barley

Application Rate

11 - 15 fl oz/A

Mode-of-Action Group

- | | |
|----|---------------|
| 27 | Pyrasulfotole |
| 6 | Bromoxynil |

Recommended Spray Volume

10-15 gal/A*

*For aerial application minimum spray volume of 5 gals/A.

/// CROP-ROTATION GUIDELINES

- The following crops have been field-tested and may be planted at the prescribed interval after an application of Huskie Complete. Huskie Complete breakdown in the soil is due mainly to microbial action. Do not plant any rotational crop within 90 days following application.

Broadleaf Weeds Controlled	
- Bittercress - Buckwheat - Cocklebur - Cockle - Cowcockle - Dandelion (seedling) - Fiddleneck - Field pennycress - Flixweed - Fumitory - Gromwell, corn - Hawksbeard - Hempnettle - Kochia - Jacob's ladder - Lambsquarters - London rocket - Marshelder - Mustard	- Nightshade* - Palmer Amaranth - Pennsylvania smartweed - Pigweed - Prickly Lettuce - Puncturevine - Radish - Ragweed - Shepherd's-purse - Smartweed - Sowthistle - Tansymustard - Velvetleaf - Volunteer Canola - Volunteer Soybean - Wallflower - Waterhemp - Wormood, (seedling)

*Controlled in spring wheat, winter, wheat, durum and Barley

7 Days	Wheat, Barley, Grain sorghum (to include grain and forage)
1 Month	Fine fescue, Tall fescue, Kentucky bluegrass, Oats, Orchardgrass, Perennial ryegrass, Annual ryegrass, Rye, Triticale
4 Months	Alfalfa ¹ , Corn, Millet ² , Soybeans
9 Months	Canola, Canaryseed, Chickpeas, Dry beans, Flax, Field Peas ³ , Green Beans, Green Peas, Lentils ⁴ , Mustards, Onions ⁵ , Potatoes, Safflower ² , Sunflowers, Sugarbeets, Timothy

¹Through tillage prior to planting alfalfa and a minimum of 12 inches of rainfall, overhead, flood irrigation, or any combination of these water sources totaling 12 inches is required between the time following a Huskie herbicide application and the time of alfalfa seeding.

²Millet and Safflower MT only, 8 inches of cumulative precipitation is required from application before planting millet or safflower in addition to the required rotational interval given in months in MT.

³Field peas: 9 months for all states except 18 months in MT.

⁴Lentils: 9 months for all states except 18 months in MN, MT, ND and SD.

⁵Onion plantback interval of 9 months is only allowed if the preceeding crop is grown with supplemental irrigation and onions are also being grown under irrigated conditions.

Where a crop is not specified, conduct a field bioassay as described in "FIELD BIOASSAY" section of this label.



LEARN MORE

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

*For more information on herbicide group numbers and mode-of-action options, visit IWillTakeAction.com

Huskie®

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Bayer, Bayer Cross and Huskie® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2025 Bayer Group. All rights reserved.

