



BROMOXYNIL	GROUP	6	HERBICIDE
PYRASULFOTOLE	GROUP	27	HERBICIDE

FIFRA Section 24(c) Special Local Need Label
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEW YORK
(EXCEPT NASSAU & SUFFOLK COUNTIES)

For Use Only By Certified Applicators
For Control of Horseweed/Marestail in Wheat, Barley, and Rye Triticale

Huskie® Herbicide

EPA Reg. 264-1023

SLN. No. NY-180001

THIS LABEL IS VALID UNTIL DECEMBER 31, 2027 UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

ENVIRONMENTAL HAZARDS

Reporting Ecological Incidents

To report ecological incidents, including mortality, injury, or harm to plants and animals, call [1-866-99BAYER (1-866-992-2937)].

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Endangered Species Protection Requirements: It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species and certain threatened species, under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product.

WEEDS CONTROLLED – WINTER AND SPRING CEREALS (WHEAT, BARLEY, RYE TRITICALE)
Apply by Ground Application Only

Partial Control			
Weed Species	Scientific name	Weed Size	Dose Rate ²
Horseweed/Marestail ¹	<i>Conyza canadensis</i>	1-4"	11-15 fl oz/A

¹ Includes ALS, phenoxy or glyphosate resistant biotypes.

²In winter wheat, 13.5 fl oz/A and an additional herbicide tank mix partner may be necessary to strengthen weed control.

Partially controlled weeds will be stunted in growth and/or be reduced in number as compared to non-treated areas and performance may not be commercially acceptable. The degree of weed control will vary with weed size, density, application coverage and growing conditions.

BROMOXYNIL	GROUP	6	HERBICIDE
PYRASULFOTOLE	GROUP	27	HERBICIDE

PRECAUTIONS

- Rainfall within 1 hour may result in reduced weed control.
- Tank mix applications of herbicides with fungicides may cause temporary yellowing, leaf burn and or height reduction of the crop.
- For optimal weed control, apply to actively growing weeds. Weed control may be reduced when weeds are under stress due to severe weather conditions, drought, very cold temperatures, etc., or under dry, dusty conditions – especially in the wheel track areas.

RESTRICTIONS FOR USE IN CEREALS

- Do not apply to crops undersown with legume species.
- Do not make more than one application of Huskie Herbicide per season.
- Do not use less than 11 fl oz/A of Huskie Herbicide unless directed by a Bayer CropScience representative.
- Do not apply more than 15 fl oz/A of Huskie Herbicide (0.037 lb pyrasulfotole/A, 0.205 lb bromoxynil/A) per season.
- Do not apply more than 0.053 pounds of mefenpyr-diethyl per acre per year.
- Do not apply Huskie Herbicide through sprinkler irrigation systems.
- Do not apply Huskie Herbicide in tank mixture with tebuconazole.
- Do not graze or harvest forage within 25 days, grain and straw within 60 days after application.
- Do not exceed 10 mph for ground application.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

**Bayer CropScience LP
800 N.Lindbergh Blvd.
St. Louis, MO 63167**