



IMIDACLOPRID	GROUP	3	INSECTICIDE
BETA-CYFLUTHRIN	GROUP	4A	INSECTICIDE

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

FIFRA Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS

For Control of Asian Citrus Psyllid on Citrus

LEVERAGE[®] 360 Insecticide

EPA Reg. No. 264-1104

SLN. No. TX-110001

THIS LABEL IS VALID UNTIL DECEMBER 31, 2025, UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED

ACTIVE INGREDIENT:

Imidacloprid*	21.0%
β-cyfluthrin**	10.5%
OTHER INGREDIENTS:	68.5%
TOTAL:	100.0%

Contains 2 lbs imidacloprid per gallon plus 1 lb β-cyfluthrin per gallon

* CAS# - 138261-41-3

** CAS# - 68359-37-5

DIRECTIONS FOR USE

Restricted Use Pesticide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

TREE and VINE CROPS
Applications for LEVERAGE 360 Insecticide

Apply specified rate per acre as a broadcast or directed foliar spray to an infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve pest control. A spray adjuvant may be used to improve coverage. LEVERAGE 360 Insecticide may be tank mixed with other insecticides for improved control of other pests.

Application rates within this label are based on full-size mature trees and vines. Use higher listed rates for moderate to heavy insect pressure. Lower rates are generally adequate for smaller trees/vines or low to moderate insect pressure but require careful scouting and may require more frequent application.

CITRUS

Crops of Crop Group 10 Including: Calamondin, Citrus citron, Citrus hybrids (*Citrus* spp., includes chironja, tangelo and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sweet and sour), Pummelo, Satsuma mandarin, and other cultivars and/or hybrids of these

Pests Controlled	Product Rate fluid ounces/Acre
Asian citrus psyllid	6.4

RESTRICTIONS

Pre-Harvest Interval (PHI): **0 day**

Minimum interval between applications: **10 days**

Maximum LEVERAGE 360 Insecticide allowed per year: **6.4 fluid ounces/Acre** (0.05 lb A/A β -cyfluthrin, 0.10 lb A/A imidacloprid)

Maximum β -cyfluthrin allowed in all forms per crop season: 0.05 lb A/A

Maximum cyfluthrin allowed in all forms per crop season: 0.10 lb A/A

Maximum β -cyfluthrin plus cyfluthrin allowed in all forms per crop season: 0.10 lb A/A

Do not apply during bloom or within 10 days prior to bloom or when bees are foraging.

LEVERAGE 360 Insecticide is to be applied through properly calibrated ground or aerial application equipment.

Minimum application volume (water): 25 GPA – ground, 25 GPA – aerial application.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)