



Luna® Flex Fungicide

Group 7 | 3 Fungicide - Bushberries

Luna® Flex Fungicide strategically counters diseases and helps reduce the severity of common problematic diseases like Scab, Melanose and Greasy Spot. The options labeled for use in a variety of crops allows for flexibility against secondary diseases like Fly Speck and Brown Rot. This makes Luna Flex a good solution for multi-crop farms to deliver high quality fruit that helps command extended performance.


/// BENEFITS

- **Broad Spectrum Control**
Labeled for key crops and diseases in the Eastern US markets
- **Great Curative Control**
Enhanced curative properties that will help improve overall disease control
- **Long-Lasting Protection**
Will provide extended performance over the recommended spray interval



/// LUNA® FLEX FUNGICIDE QUICK FACTS

- **Active Ingredients:** Fluopyram and Difenoconazole
- **FRAC:** Group 7 and 3 fungicide
- **Formulation Type:** Liquid, 375 SC (Soluble Concentrate)
- **Signal Word:** Caution
- **REI:** 12 hours
- **PHI:** 7 days
- **Use Rate:** 11.2 - 13.6 oz./A

Key Bushberry Diseases	Additional Registered Crops
• Alternaria • Anthracnose • Gray Mold • Leaf Rust • Monilinia Blight • Mummy Berry • Powdery Mildew	• Apples • Bulb Vegetables • Citrus • Cucurbits • Fruiting Vegetables • Pome Fruits • Stone Fruits • Strawberries



Luna® Flex Fungicide Effectively Controls Monilinia blight and Mummy berry (Monilinia spp.)



Wild blueberry trial conducted in Nunica, MI in 2021 by Michigan State University. Treatments were applied two times: May 11 (late green tip) and May 18 (pink bud). Other fungicide applications in the programs were similar across all the treatments. Nonionic surfactant was used with all spray applications. Data for shoot strikes and infected fruits were recorded on May 29 and July 10, respectively. Means within category followed by the same letter are not significantly different according to Fisher's Protected LSD ($P < 0.05$)

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

Luna Flex contains oil in the formulation. If tank-mixing Luna Flex with oil-sensitive products, read and follow their label restrictions. Do not apply in tank-mix with, or 7 days before or after an application of Captan.

For more information, visit www.CropScience.Bayer.us.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Bayer, Bayer Cross and Luna® Flex are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2025 Bayer Group. All rights reserved.

**Luna®
FLEX**

