



Luna® Flex Fungicide

Group 7 / 3 Fungicide - Stone Fruits

Luna® Flex Fungicide strategically counters diseases and helps reduce the severity of common problematic diseases like Scab, Melanose and Greasy Spot. The options labeled for use in a variety of crops allows for flexibility against secondary diseases like Fly Speck and Brown Rot. This makes Luna Flex a good solution for multi-crop farms to deliver high quality fruit that helps command extended performance.

/// BENEFITS

- **Broad Spectrum Control**

Labeled for key crops and diseases in the Eastern US markets

- **Great Curative Control**

Enhanced curative properties that will help improve overall disease control

- **Long-Lasting Protection**

Will provide extended performance over the recommended spray interval

/// LUNA® FLEX FUNGICIDE QUICK FACTS

- **Active Ingredients:** Fluopyram and Difenconazole
- **FRAC:** Group 7 and 3 fungicide
- **Formulation Type:** Liquid, 375 SC (Soluble Concentrate)
- **Signal Word:** Caution
- **REI:** 12 hours
- **PHI:** 0 days
- **Use Rate:** 11.0 - 13.6 oz./A

Luna® Flex Fungicide Effectively Controls Cherry Leaf Spot and Helps Retain Foliage Late in the Season.




Cherry Leaf Spot
Untreated Control vs Luna® Flex Fungicide

Picture from trial conducted at MI in 2021. Luan Flex was applied in April-May; picture taken on September 29, 2021

Labeled Stone Fruit Diseases	Additional Registered Crops
• Alternaria Spot and Fruit Rot • Anthracnose • Brown Rot Blossom Blight Fruit Rot • Cherry Leaf Spot • Green Fruit Rot • Jacket Rot • Powdery Mildew • Ripe Fruit Rot • Rusty Spot • Rust • Scab • Shot hole	• Apples • Bulb Vegetables • Bushberries • Citrus • Cucurbits • Fruiting Vegetables • Pome Fruits • Strawberries


Luna® Flex Fungicide Effectively Controls Cherry Leaf Spot



% of Cherry Leaf Spot

Tart Cherry (cv. Montmorency). Trial conducted in Lyons, NY in 2021 by an independent research organization. Treatments were applied on June 11 and August 7. Other fungicide applications in the programs were similar across all the treatments. Luna Flex was applied at a lower rate for testing purpose. Follow the label for rate and timings for commercial applications. Nonionic surfactant was used with all spray applications. Data recorded on October 21. Means within category followed by the same letter are not significantly different according to Fisher's Protected LSD ($P < 0.05$).


Luna® Flex Provides Effective Control of Cherry Leaf Spot and Powdery Mildew



■ Powdery Mildew ■ Cherry Leaf Spot

Tart Cherry (cv. Montmorency). Trial conducted in East Lansing, MI in 2021 by Michigan State University. Treatments were applied on June 1 and June 24. Other fungicide applications in the programs were similar across all the treatments. Luna Flex was applied at a lower rate for testing purpose. Follow the label for rate and timings for commercial applications. Nonionic surfactant was used with all spray applications. Data recorded on July 22 and August 22. Means within category followed by the same letter are not significantly different according to Fisher's Protected LSD ($P < 0.05$).

Peach Fruit Rot Control: Luna® Flex vs. Competitors



■ 4 days after harvest ■ 7 days after harvest

Peach (cv. Flame Prince). Trial conducted in Byron, GA in 2021 by University of Georgia. Treatments were applied as preharvest sprays on July 19 and July 26. Luna Flex was applied at a lower rate for testing purpose. Follow the label for rate and timings for commercial applications. Please follow the label for rates and timings for commercial applications. Nonionic surfactant was used with all spray applications. Brown rot incidence was recorded on fruit stored at ambient temperature. Means within category followed by the same letter are not significantly different according to Fisher's Protected LSD ($P < 0.05$).

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

Luna Flex contains oil in the formulation. If tank-mixing Luna Flex with oil-sensitive products, read and follow their label restrictions. Do not apply in tank-mix with, or 7 days before or after an application of Captan.

For more information, visit www.CropScience.Bayer.us.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Bayer, Bayer Cross and Luna® Flex are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2025 Bayer Group. All rights reserved.

**Luna®
FLEX**

