



Luna[®]
FLEX

Luna[®] Flex Fungicide

Group 7 | 3 Fungicide - Strawberries

Luna[®] Flex Fungicide strategically counters diseases and helps reduce the severity of common problematic diseases like Scab, Melanose and Greasy Spot. The options labeled for use in a variety of crops allows for flexibility against secondary diseases like Fly Speck and Brown Rot. This makes Luna Flex a good solution for multi-crop farms to deliver high quality fruit that helps command extended performance.


/// BENEFITS

- **Broad Spectrum Control**
Labeled for key crops and diseases in the Eastern US markets
- **Great Curative Control**
Enhanced curative properties to improve overall disease control
- **Long-Lasting Protection**
Extended performance over the recommended spray interval


/// LUNA[®] FLEX FUNGICIDE QUICK FACTS

- **Active Ingredient:** Fluopyram and Difenoconazole
- **FRAC:** Group 7 and 3 fungicide
- **Formulation Type:** Liquid, 375 SC (Soluble Concentrate)
- **Signal Word:** Caution
- **REI:** 12 hours
- **PHI:** 0 days
- **Use Rate:** 12.0 - 13.6 fl. oz./A

Key Strawberry Diseases	Additional Registered Crops
• Anthracnose • Gray Mold • Leaf Rust • Leaf Spot • Powdery Mildew • Rhizopus Fruit Rot	• Apples • Bulb Vegetables • Bushberries • Citrus • Cucurbits • Fruiting Vegetables • Pome Fruits • Stone Fruits



Luna® Flex Fungicide Effectively Controls Strawberry Botrytis Fruit Rot



Trial conducted in Plant City, FL in 2021 by University of Florida. Test products were applied when weather conditions for Botrytis cinerea infection were favorable (17 to 25°C and ° 12 h leaf wetness) according to the Strawberry Advisory System. For head-to-head comparison test products were applied three times. Please follow the label for rates and timings for commercial applications. Other fungicide applications in the programs were similar across all the treatments. Yield was recorded based on the weight of marketable fruit and Botrytis fruit rot incidence was recorded as the percentage of symptomatic fruit relative to all harvested fruit. Means within series followed by the same letter are not significantly different according to Fisher's Protected LSD ($P < 0.05$).

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

Luna Flex contains oil in the formulation. If tank-mixing Luna Flex with oil-sensitive products, read and follow their label restrictions. Do not apply in tank-mix with, or 7 days before or after an application of Captan.

For more information, visit www.CropScience.Bayer.us.

**Luna®
FLEX**

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Bayer, Bayer Cross and Luna® Flex are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2025 Bayer Group. All rights reserved.

