



Luna® Pro Fungicide

Group 7 / 3 Fungicide

Luna® Pro is a new premix fungicide from Bayer that provides foliar and soilborne disease control in potatoes. It is a broad-spectrum solution for potato diseases, with the performance you've come to expect from the Luna® family of fungicides.

/// BENEFITS

- **Consistent Results**
Consistently strong disease control from two proven active ingredients, providing an effective combination of preventative and curative activity.
- **Systemic Activity**
Unique combination of systemic active ingredients provide protection throughout the plant, protecting against foliar and soilborne pathogens.
- **Resistance Management**
The first DMI-containing foliar fungicide in potatoes by Bayer, providing rotational flexibility to help manage resistance development.
- **Enhanced White Mold Control**
Provides exceptional white mold control while also protecting against other key foliar diseases.
- **High Level of Yield Potential**
Contains active ingredients with proven yield enhancement.

/// LUNA® PRO FUNGICIDE QUICK FACTS


- **Active Ingredient:** Fluopyram and Prothioconazole
- **FRAC:** Group 7 and 3 fungicide
- **Formulation Type:** Liquid, 375 SC (Soluble Concentrate)
- **REI:** 12 hours
- **PHI:** 7 days
- **Use Rate:** 10 oz./A

Key Diseases

- Black Dot
- Botrytis Leaf Spot
- Brown Spot
- Early Blight
- White Mold




Trials Show that Luna® Pro Fungicide is Highly Effective Against Early Blight and Helps Realize Yield Potential



Average of data from 14 trials conducted from 2017-2021 in ID, WA, ND, CO, MN, WI, MI, MN


Data shows that Luna® Pro Fungicide is highly effective on White Mold

Miller Research - Rupert, ID
White Mold



Trials conducted in ID in 2019 and 2020

Trials Show that Luna® Pro Fungicide is Highly Effective Against Early Blight and Helps Realize Yield Potential when Used in a Program



Trial conducted in Rupert, ID in 2021 by an independent research firm. Treatments were applied at row closure and 14 days after row closure. Bravo @ 1.0 pt/A was applied as tank mix to all treatments. AUDPC = Area Under Disease Progress Curve. Means within category followed by the same letter are not significantly different according to Fisher's Protected LSD (P<0.05)

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

For more information, visit www.CropScience.Bayer.us.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Performance may vary, from location to location and from year to year, as local growing, soil and environmental conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on their growing environment. The recommendations in this material are based upon trial observations and feedback received from a limited number of growers and growing environments. These recommendations should be considered as one reference point and should not be substituted for the professional opinion of agronomists, entomologists or other relevant experts evaluating specific conditions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your state department(s) of agriculture, local dealer or representative for the product registration status in your state. Not all products are registered for use in all states and may be subject to use restrictions. Bayer, Bayer Cross, Luna Tranquility® and Luna® Pro Fungicide are registered trademarks of Bayer Group. All other trademarks are the property of their respective owners. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LLC, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2026 Bayer Group. All rights reserved.

**Luna®
PRO**

