



Luna Sensation® Fungicide

Group 7 / 11 Fungicide - Lettuce

Luna Sensation® fungicide protects against key diseases throughout the growing season, resulting in improved plant health as well as high-quality crops, and provides an effective approach to resistance management.

/// BENEFITS

Exceptional Efficacy

Luna Sensation® fungicide consistently provides excellent disease control of Lettuce drop and Powdery mildew.

Resistance Management

Luna Sensation contributes to resistance management by containing two modes of action as well as offering an excellent rotational fit with other available fungicides.

Yield Potential

Luna Sensation applied in season and/or prior to harvest helps improve quality at harvest as well as through transit and storage.

/// LUNA SENSATION® FUNGICIDE QUICK FACTS

Active Ingredients: Fluopyram and Trifloxystrobin

FRAC: Group 7 and 11 Fungicide

Signal Word: Caution

REI: 12 hours

Use Rate: 7.6 oz/A


PHI: 0 day for broadcast foliar uses; 20 day for banded applications

Approved for aerial applications.

Labeled Diseases	
Control	Suppression
• Alternaria leaf spot • Powdery mildew • Lettuce drop (<i>Sclerotinia</i> spp.)	• Rhizoctonia bottom rot




Luna Sensation® Fungicide Application Supports High Yields in Lettuce




Year: 2016
Location: Chualar, CA
Single trial - Application made at thinning (4-6 leaf stage) for control of *Sclerotinia minor* 30 days after planting
Percent marketable heads counted at harvest 57 days after planting

Luna Sensation® Fungicide Shows Strong Yield Potential even under Heavy Disease Pressure



Year: 2016
Location: King City, CA
Single trial - Application made at thinning (4-6 leaf stage) for control of *Sclerotinia* (heavy disease pressure)
Percent marketable heads counted at harvest, 45 days after application

Excellent Control of *Sclerotinia minor*



Year: 2017
Location: Aromas, CA
Single trial - First application made at thinning (4-6 leaf stage) and second application made 14 days later for control of *Sclerotinia minor* Percent disease incidence at 66 and 76 days after planting

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

For more information, visit www.CropScience.Bayer.us.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Performance may vary, from location to location and from year to year, as local growing, soil and environmental conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on their growing environment. The recommendations in this material are based upon trial observations and feedback received from a limited number of growers and growing environments. These recommendations should be considered as one reference point and should not be substituted for the professional opinion of agronomists, entomologists or other relevant experts evaluating specific conditions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your state department(s) of agriculture, local dealer or representative for the product registration status in your state. Not all products are registered for use in all states and may be subject to use restrictions. Bayer, Bayer Cross and Luna Sensation® are registered trademarks of Bayer Group. All other trademarks are the property of their respective owners. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LLC, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2026 Bayer Group. All rights reserved.

**Luna
SENSATION®**

