



# Luna EXPERIENCE®

Net Contents:

1/2 Gal. (64 fl. oz.)

FLUOPYRAM	GROUP 7	FUNGICIDE
TEBUCONAZOLE	GROUP 3	FUNGICIDE

**Broad spectrum fungicide for control of plant diseases.**

<b>ACTIVE INGREDIENTS:</b> FLUOPYRAM*	17.6%
TEBUCONAZOLE*	17.6%
<b>OTHER INGREDIENTS:</b>	64.8%
<b>TOTAL: 100.0%</b>	

Contains 1.67 lbs FLUOPYRAM and 1.67 lbs TEBUCONAZOLE per gallon  
\*(CAS Numbers 658066-35-4 and 107534-96-3)

EPA Reg. No. 264-1091

SUSPENSION CONCENTRATE

## KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements and directions for use on label.

Produced for:

Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167

Luna Experience is a registered trademark of Bayer Group.

©2022 Bayer CropScience

07/22

210915D

US86247194D

**Almond; Brassica leafy greens (subgroup 5B); Bulb vegetables (group 3-07); Cucurbits (group 9); Grapes; Hops; Dry and Succulent Beans; Okra; Pistachio; Stone Fruits (group 12); Sunflower; Tree Nuts (group 14).**

### FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

**For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.**

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.

Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

**NOTE TO PHYSICIAN:** Treat Symptomatically.

**For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

#### CAUTION

Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

### **Personal Protective Equipment (PPE):**

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant (such as nitrile or butyl) gloves.

### **User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **Engineering Control Statements**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

## **USER SAFETY RECOMMENDATIONS**

### **Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## **ENVIRONMENTAL HAZARDS**

For terrestrial uses: This pesticide is toxic to mammals, fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when disposing of equipment wash water or rinsate. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of Fluopyram. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

#### **DIRECTIONS FOR USE**

#### **STOP - READ THE LABEL BEFORE USE**

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

**Read entire label before using this product.**

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Not for sale, distribution, or use in Nassau and Suffolk counties, New York except as permitted under FIFRA 24(c), Special Local Need registration.**

**Aerial Application Prohibited in New York State.**

#### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 12 hours for all crops except when performing cane tying, turning, or girdling on grapes.**

**The REI is 5 days for treated grapes when conducting cane tying, turning, or girdling on grapes. The REI is 12 hours for all other activities in grapes.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls over long-sleeved shirt and long pants
- socks and shoes
- chemical-resistant gloves made of any waterproof material such as natural rubber  $\geq$  14 mils

## **PRODUCT INFORMATION**

LUNA EXPERIENCE<sup>®</sup> is a broad-spectrum fungicide with preventative, systemic and curative properties labeled for the control or suppression of certain crop diseases.

### **LABELED USES**

Almond; Brassica leafy greens (subgroup 5B); Bulb vegetables (group 3-07); Cucurbits (group 9); Grapes; Hops; Dry and Succulent Beans; Okra; Pistachio; Stone Fruits (group 12); Sunflower; Tree Nuts (group 14).

### **RESISTANCE-MANAGEMENT RECOMMENDATIONS**

For resistance management, please note that LUNA EXPERIENCE contains both a Group 7 and Group 3 fungicide. Any fungal population may contain individuals naturally resistant to LUNA EXPERIENCE and other Group 7 or Group 3 fungicides. A gradual or total loss of pest control may occur over time if these (fungicides) are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of LUNA EXPERIENCE or other Group 7 or Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

## **MANDATORY SPRAY DRIFT**

### **Aerial Applications**

- For aerial applications, **DO NOT** apply when wind speeds exceed 10 mph at the application site.
- For all applications, applicators are required to use a medium or coarser droplet size (ASABE S572. I).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

### **Airblast applications**

- Sprays must be directed into the canopy.
- For all applications, applicators are required to use a medium or coarser droplet size (ASABE S572. I).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions

### **Ground Boom Applications**

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser droplet size (ASABE S572. I).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

## **SPRAY DRIFT ADVISORIES**

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

#### **Controlling Droplet Size - Ground Boom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

#### **Controlling Droplet Size – Aircraft**

- Adjust Nozzles - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### **BOOM HEIGHT – Ground Boom**

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **RELEASE HEIGHT – Aircraft**

Higher release heights increase the potential for spray drift. When applying aerially to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

## **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

## **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversion.

## **WIND**

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## **BOOM-LESS GROUND APPLICATIONS**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

## **HANDHELD TECHNOLOGY APPLICATIONS**

Take precautions to minimize spray drift.

## **APPLICATION INFORMATION**

Use sufficient water volume to provide thorough and uniform coverage to obtain the most effective disease control.

### **Ground Application**

For optimum disease control, apply in sufficient water to ensure thorough coverage of foliage, bloom, and fruit.

### **Aerial Application**

For aerial application equipment, a minimum of **10** gallons of water per acre for tree crops and **2** gallons of water per acre for field and vegetable crops is required. Aerial Application Prohibited in New York State.



### **Air-Blast Application**

Air-assisted or air-blast sprayers move spray droplets into the crop canopy using a forced-air system. The fan should be set up to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area. Equip sprayers with nozzles that provide accurate and uniform application.

### **Chemigation Application**

Apply this product only through center pivot, motorized-lateral move, traveling gun, solid set or portable (wheel move, side roll, end tow, or hand move) irrigation systems. **DO NOT** apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, must shut the system down and make necessary adjustments should the need arise.

LUNA EXPERIENCE has not been sufficiently tested when applied through irrigation systems to assure consistent product performance for all labeled uses. The following application techniques are provided for user reference but do not constitute a warranty of fitness for application through sprinkler irrigation equipment.

**DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label prescribed safety devices for public water systems are in place. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an alternative to the RPZ, the water from the public water system must be discharged into a reservoir tank prior to pesticide introduction.

There must be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. Pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and

connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The systems must contain functional interlocking controls, to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

**DO NOT** apply when wind speed favors drift. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Apply pesticide continuously for the duration of the water application. For mixing instructions, please refer to specific use directions in the "Spray mixing and compatibility" section.

This product may be used through two basic types of irrigation systems as outlined in **Sections A and B** below. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Determine which type of irrigation system is in place, then refer to the appropriate directions provided below for each type. See crops section on the label for required treatment rates and additional use information.

### **A. Center Pivot, Motorized-Lateral Move and Traveling Gun Irrigation Equipment**

For injections of pesticides, these continuously moving systems must use a positive displacement injection pump of either diaphragm or piston type and be constructed of materials that are compatible with pesticides. They must also be capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems. Thoroughly mix required amount of this product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until this product has been cleared from the last sprinkler head.

## **B. Solid-Set, Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Equipment**

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred to support even and quick distribution. However, a positive-displacement pump can also be used. For solid set systems, determine acreage covered by sprinkler. Fill the tank of injection equipment with water and adjust flow to use contents over 30 to 45 minutes. Mix desired amount of this product for acreage to be covered with water so that the total mixture of this product plus water in the injection tank is equal to the quantity of water used during calibration. Provide chemical supply tank agitation sufficient for mixing until chemigation is completed. Operate entire system at normal pressures recommended by the manufacturer of injection equipment used, for amount of time established during calibration. This product can be injected during the irrigation cycle or as a separate application. Stop injection equipment with any system after treatment is completed and continue to operate irrigation system until this product has been cleared from the last sprinkler head.

## **SPRAY MIXING AND COMPATIBILITY**

Begin with clean spray equipment and add one-half of the required amount of water to the spray or mixing tank and start agitation. Add the required quantity of fungicide and the tank-mix partner if applicable to the water and complete filling with water to the required total volume. Follow the recommendations of your State Cooperative Extension Service for tank-mixing with other products. In general, follow the order beginning first with water conditioners, water soluble packaging (wait for it to completely dissolve), wettable powders and water-dispersible granular products, liquid flowables and suspension concentrates, emulsifiable concentrates, and adjuvants last. Maintain agitation throughout spraying. **DO NOT** allow spray mixture to remain in the tank overnight, or for long periods during the day without agitation.

LUNA EXPERIENCE is physically compatible with most commonly used fungicide, herbicide, insecticide, and foliar nutrient products. However, the compatibility of LUNA EXPERIENCE with all potential tank-mix partners has not been fully investigated. If tank mixing with other pesticides is desirable, conduct a jar test with the volumes and rates typically used in agricultural application. Using a small container of water, add the proportionate amounts of the products: wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other

precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

The crop safety of all potential tank-mixes with LUNA EXPERIENCE has not been tested on all crops listed on the label. Before applying any tank-mixture not specified on this label, safety to the target crop must be confirmed on a small portion of the crop listed on the label to be treated to ensure an adverse response will not occur.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### **PRODUCT RESTRICTIONS AND LIMITATIONS**

**DO NOT** apply more than the maximum yearly rate for each specific crop from any combination of products containing FLUOPYRAM or TEBUCONAZOLE.

### **ROTATIONAL CROP RESTRICTIONS**

Areas treated with this product can be replanted immediately following harvest with any crop for which both a FLUOPYRAM and a TEBUCONAZOLE tolerance exist. This includes: Almond; Barley; Brassica leafy greens (subgroup 5B); Bulb vegetables (group 3-07); Corn, field, grain; Corn, pop, grain; Corn, sweet; Cotton; Cucurbits (group 9); Dry and Succulent beans; Fruiting Vegetables (group 8-10); Grapes; Hops; Peanut; Pecan; Pistachio; Pome fruit (group 11); Soybean; Stone Fruits (group 12); Sunflower; Tree Nuts (group 14) and Wheat.

The following crops can be planted into treated areas 120 days after the last application of LUNA EXPERIENCE:

Alfalfa; Artichoke,(Globe); Brassica leafy greens subgroup 5A; Carrot; Cereals (except corn and rice) (group 15) which include: buckwheat, millet (pearl and proso), oats, rye, sorghum, teosinte, and triticale; Citrus (group 10-10); Dill seed; Ginseng; Herb (subgroup 19A); Leafy vegetables (except watercress) (group 4); Potato and other root, tuberous and corm vegetables (except sugarbeet) (subgroups 1B and 1C); Rapeseed (subgroup 20A); Small Berries (caneberries and bushberries ) (subgroups 13-07A and 13-07B); Strawberry and other low-growing berries, except cranberry (subgroup 13-07G); Succulent Shelled Pea and Bean subgroup 6B; Sugarcane (in Region 3); and Sugarbeet.

**DO NOT** rotate to crops other than those listed above.

## USE DIRECTIONS FOR SPECIFIC CROPS

### ALMOND

Disease Control	Application Rate	Product Instructions
Brown rot blossom blight ( <i>Monilinia laxa</i> ) ( <i>Monilinia fructicola</i> )	6.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Shot hole ( <i>Wilsonomyces carpophilus</i> )	(0.078 – 0.222 lb/acre fluopyram)	
Powdery mildew ( <i>Sphaerotheca pannosa</i> ) ( <i>Podosphaera tridactyla</i> )	(0.078 – 0.222 lb/acre tebuconazole)	
Jacket rot <sup>1</sup>		
Green fruit rot ( <i>Botrytis cinerea</i> )		
Alternaria leaf spot <sup>1</sup> ( <i>Alternaria alternata</i> )		
Anthracnose <sup>1</sup> ( <i>Colletotrichum acutatum</i> )		
Scab <sup>1</sup> ( <i>Cladosporium carpophilum</i> )		
Rust <sup>1</sup> ( <i>Tranzschelia discolor</i> )		

**Restrictions:**

- Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 7 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 35 days of harvest.
- **DO NOT** cut cover crops in treated areas for feed, or allow livestock to graze treated areas.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

<sup>1</sup>Not for use in CA without a supplemental label.

**BRASSICA LEAFY GREENS (Subgroup 5B)**

Broccoli Raab; Cabbage, Chinese (Bok Choy); Collards; Kale; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens.

Disease Control	Application Rate	Product Instructions
Powdery mildew <sup>1</sup> ( <i>Erysiphe cruciferarum</i> ) ( <i>Erysiphe polygoni</i> )	6.0 to 8.6 fl oz/acre (0.078 – 0.112 lb/acre fluopyram)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Cercospora leaf spot <sup>1</sup> ( <i>Cercospora brassicicola</i> )		
Alternaria blight / Black spot <sup>1</sup> ( <i>Alternaria</i> spp.)	(0.078 – 0.112 lb/acre tebuconazole)	

Botrytis gray mold ( <i>Botrytis cinerea</i> ) Sclerotinia stem rot ( <i>Sclerotinia sclerotiorum</i> ) ( <i>Sclerotinia minor</i> )	8.6 fl oz/acre (0.112 lb/acre fluopyram) (0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the shorter intervals.
--	--	---

Disease Suppression	Application Rate	Product Instructions
Rhizoctonia bottom rot ( <i>Rhizoctonia solani</i> )	8.6 fl oz/acre (0.112 lb/acre fluopyram) (0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the shorter intervals.

#### Restrictions:

- Maximum single application rate: 8.6 fl oz/acre of LUNA EXPERIENCE (0.112 lb/acre fluopyram and 0.112 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 7 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.450 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- Application to Turnip greens is limited to east of the Rockies.

<sup>1</sup>Not for use in CA without a supplemental label.

## BULB VEGETABLES (group 3-07)

**Onion, bulb subgroup:** Daylily, Bulb; Fritillaria, Bulb; Garlic, Bulb; Garlic, Great-Headed, Bulb; Garlic, Serpent, Bulb; Lily, Bulb; Onion, Bulb; Onion, Chinese, Bulb; Onion, Pearl; Onion, Potato, Bulb; Shallot, Bulb; cultivars, varieties, and/or hybrids of these.

**Onion, green subgroup:** Chive, Fresh Leaves; Chive, Chinese, Fresh Leaves; Elegans Hosta; Fritillaria, Leaves; Kurrat; Lady's Leek; Leek; Leek, Wild; Onion, Beltsville Bunching; Onion, Fresh; Onion, Green; Onion, Macrostem; Onion, Tree, Tops; Onion, Welsh, Tops; Shallot, Fresh Leaves; cultivars, varieties, and/or hybrids of these.

Disease Control	Application Rate	Product Instructions
Botrytis leaf blight, neck rot ( <i>Botrytis squamosa</i> ) ( <i>Botrytis allii</i> ) Purple blotch ( <i>Alternaria porri</i> ) Rust ( <i>Puccinia allii</i> ) ( <i>Puccinia porri</i> )	8.0 to 12.8 fl oz/acre (0.104 – 0.167 lb/acre fluopyram)  (0.104 – 0.167 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Stemphylium leaf blight / stalk rot ( <i>Stemphylium vesicarium</i> ) Cladosporium leaf blotch ( <i>Cladosporium allii</i> )	12.8 fl oz/acre (0.167 lb/acre fluopyram) (0.167 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Disease Suppression	Application Rate	Product Instructions
White rot ( <i>Sclerotium cepivorum</i> )	12.8 fl oz/acre (0.167 lb/acre fluopyram) (0.167 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.



**Restrictions:**

- Maximum single application rate: 12.8 fl oz/acre of LUNA EXPERIENCE (0.167 lb/acre fluopyram and 0.167 lb/acre tebuconazole).
- Maximum annual application rate: 25.6 fl oz/acre of LUNA EXPERIENCE (0.334 lb/acre fluopyram and 0.334 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 3 (at 8.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 10 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.335 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

**CUCURBITS (Group 9)**

**Melon subgroup:** Citron Melon; Muskmelon; Watermelon.

**Squash/Cucumber subgroup:** Chayote (Fruit); Chinese Waxgourd; Cucumber; Gherkin; Gourd, Edible; *Momordica* spp.; Pumpkin; Squash, Summer; Squash, Winter.

Disease Control	Application Rate	Product Instructions
Powdery mildew ( <i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i> ) ( <i>Erysiphe cichoracearum</i> ) Alternaria leaf spot ( <i>Alternaria cucumerina</i> )	6.0 to 17 fl oz/acre (0.078 – 0.222 lb/acre fluopyram) (0.078 – 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Gummy stem blight <sup>1</sup> ( <i>Didymella bryoniae</i> )	10.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Belly rot <sup>1</sup> ( <i>Rhizoctonia solani</i> )	(0.130 - 0.222 lb/acre fluopyram)	
Anthracnose <sup>1</sup> ( <i>Colletotrichum</i> spp.)	(0.130 - 0.222 lb/acre tebuconazole)	

**Note:** A mild yellowing on leaf margins is sometimes seen following application of LUNA EXPERIENCE in cucurbits.

**Restrictions:**

- Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 10 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.691 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

<sup>1</sup>Not for use in CA without a supplemental label.

**GRAPES**

<b>Disease Control</b>	<b>Application Rate</b>	<b>Product Instructions</b>
Powdery mildew ( <i>Uncinula necator</i> )	6.0 to 8.6 fl oz/acre (0.078 – 0.112 lb/acre fluopyram) (0.078 – 0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Botrytis bunch rot / Gray mold ( <i>Botrytis cinerea</i> ) Black rot ( <i>Guignardia bidwellii</i> ) Sour rot <sup>1</sup> ( <i>various pathogens</i> )	8.0 to 8.6 fl oz/acre (0.104 – 0.112 lb/acre fluopyram) (0.104 – 0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
<b>Disease Suppression</b>	<b>Application Rate</b>	<b>Product Instructions</b>
Phomopsis cane and leaf spot ( <i>Phomopsis viticola</i> )	8.6 fl oz/acre (0.112 lb/acre fluopyram) (0.112 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. When disease pressure is severe, use the shorter intervals.

**Restrictions:**

- Maximum single application rate: 8.6 fl oz/acre of LUNA EXPERIENCE (0.112 lb/acre fluopyram and 0.112 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 12 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE after BBCH 73 stage (when bunches begin to hang) or within 45 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

<sup>1</sup>Not for use in CA without a supplemental label.

**HOPS**

Disease Control	Application Rate	Product Instructions
Powdery mildew ( <i>Sphaerotheca humuli</i> ) ( <i>Sphaerotheca macularis</i> )	8.0 to 17 fl oz/acre  (0.104 - 0.222 lb/acre fluopyram)  (0.104 - 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.  Make applications in sufficient water volume (just to the point of runoff). Thorough coverage is needed for disease control. Apply increasing product rate as plant size increases.  8.0 fl oz with 15 - 30 gals./acre 10.0 fl oz with 31 - 60 gals./acre 12.0 fl oz with 61 - 90 gals./acre 14.0 fl oz with 91 - 200 gals./acre

Disease Suppression	Application Rate	Product Instructions
Gray mold ( <i>Botrytis</i> spp.)	17 fl oz/acre (0.222 lb/acre fluopyram) (0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.

**Restrictions:**

- Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 4 (at 8.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 14 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 14 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

**DRY AND SUCCULENT BEANS<sup>1</sup>**

Disease Control	Application Rate	Product Instructions
Powdery mildew ( <i>Erysiphe graminis</i> ) Septoria leaf and glume blotch ( <i>Septoria</i> spp.) Rusts ( <i>Puccinia</i> spp.)	8.0 to 12.8 fl oz/acre (0.104 – 0.167 lb/acre fluopyram) (0.104 – 0.167 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

**Restrictions:**

- Maximum single application rate: 12.8 fl oz/acre of LUNA EXPERIENCE (0.167 lb/acre fluopyram and 0.167 lb/acre tebuconazole).
- Maximum annual application rate: 25.6 fl oz/acre of LUNA EXPERIENCE (0.334 lb/acre fluopyram and 0.334 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 3 (at 8.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 10 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.672 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 14 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- **DO NOT** allow livestock to graze treated area for 14 days and **DO NOT** harvest for food or feed within 14 days of application.

<sup>1</sup>Not for use in CA without a supplemental label.

**OKRA**

Disease Control	Application Rate	Product Instructions
Cercospora leaf spot ( <i>Cercospora</i> spp.)	12.8 fl oz/acre (0.167 lb/acre fluopyram) (0.167 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.

**Restrictions:**

- Maximum single application rate: 12.8 fl oz/acre of LUNA EXPERIENCE (0.167 lb/acre fluopyram and 0.167 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 2
- Minimum retreatment interval: 14 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.675 lbs tebuconazole per acre per year including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 3 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

**PISTACHIO**

Disease Control	Application Rate	Product Instructions
Botryosphaeria panicle and shoot blight ( <i>Botryosphaeria dothidea</i> ) Septoria leaf spot ( <i>Septoria pistaciarum</i> ) Blossom and shoot blight ( <i>Botrytis cinerea</i> ) Alternaria late blight ( <i>Alternaria alternata</i> )	6.4 to 17 fl oz/acre (0.083 – 0.222 lb/acre fluopyram) (0.083 – 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

**Restrictions:**

- Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.4 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 10 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 35 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- **DO NOT** cut cover crops in treated areas for feed, or allow livestock to graze treated areas.

**STONE FRUITS (Group 12)**

Apricot; Cherry, sweet; Cherry, tart; Nectarine; Peach; Plum ; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plumcot; Prune (fresh).

Disease Control	Application Rate	Product Instructions
Brown rot blossom blight Fruit rot ( <i>Monilinia</i> spp.) Powdery mildew Rusty spot ( <i>Podosphaera</i> spp.) ( <i>Sphaerotheca pannosa</i> ) Shot hole <sup>1</sup> ( <i>Wilsonomyces carpophilus</i> ) Jacket rot, Green fruit rot <sup>1</sup> ( <i>Botrytis cinerea</i> ) Peach leaf curl <sup>1</sup> ( <i>Taphrina</i> spp.)	6.0 to 10 fl oz/acre (0.078 – 0.130 lb/acre fluopyram) (0.078 – 0.130 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.



Scab ( <i>Cladosporium carpophilum</i> )	8.0 to 10 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Rust ( <i>Tranzschelia discolor</i> )	(0.104 – 0.130 lb/acre fluopyram) (0.104 – 0.130 lb/acre tebuconazole)	

#### Restrictions:

- Maximum single application rate: 10 fl oz/acre of LUNA EXPERIENCE (0.130 lb/acre fluopyram and 0.130 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 5 (at 6.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 7 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 1.35 lbs tebuconazole per acre per year, including soil and foliar uses.
- May be applied the day of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

<sup>1</sup>Not for use in CA without a supplemental label.

## SUNFLOWER

Disease Control	Application Rate	Product Instructions
Rust ( <i>Puccinia helianthi</i> )	9.0 to 12.8 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. When disease pressure is severe, use the higher rates.
Alternaria leaf spot ( <i>Alternaria</i> spp.)	(0.117 – 0.167 lb/acre fluopyram)	
Sclerotinia wilt and head rot ( <i>Sclerotinia sclerotiorum</i> )	(0.117 – 0.167 lb/acre tebuconazole)	
Powdery mildew ( <i>Erysiphe cichoracearum</i> )		

**Restrictions:**

- Maximum single application rate: 12.8 fl oz/acre of LUNA EXPERIENCE (0.167 lb/acre fluopyram and 0.167 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 3 (at 9.0 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 14 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.450 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 50 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

**TREE NUTS (Group 14)***(See separate use directions for: Almonds and Pistachio)*

Beech nut; Brazil nut; Butternut; Cashew; Chestnut; Chinquapin; Filbert (hazelnut); Hickory nut; Macadamia nut (bush nut); Walnut, black and English (Persian).

<b>Disease Control</b>	<b>Application Rate</b>	<b>Product Instructions</b>
Botryosphaeria panicle and shoot blight <i>(Botryosphaeria dothidea)</i>	10.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14- to 21-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Brown rot blossom blight <i>(Monilinia laxa)</i> <i>(Monilinia fructicola)</i>	(0.130 - 0.222 lb/acre fluopyram)	
Alternaria leaf and fruit spot <i>(Alternaria alternata)</i>	(0.130 - 0.222 lb/acre tebuconazole)	
Alternaria late blight <i>(Alternaria alternata)</i>		
Scab <i>(Cladosporium carpophilum)</i> <i>(Cladosporium caryigenum)</i>		
Shothole <i>(Wilsonomyces carpophilus)</i>		
Anthracnose <sup>1</sup> <i>(Colletotrichum acutatum)</i> <i>(Glomerella cingulata)</i>		
<b>Disease Suppression</b>	<b>Application Rate</b>	<b>Product Instructions</b>
Eastern filbert blight <i>(Anisogramma anomala)</i>	10.0 to 17 fl oz/acre (0.130 - 0.222 lb/acre fluopyram) (0.130 - 0.222 lb/acre tebuconazole)	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

**Restrictions:**

- Maximum single application rate: 17 fl oz/acre of LUNA EXPERIENCE (0.222 lb/acre fluopyram and 0.222 lb/acre tebuconazole).
- Maximum annual application rate: 34 fl oz/acre of LUNA EXPERIENCE (0.444 lb/acre fluopyram and 0.444 lb/acre tebuconazole) per year.
- Maximum number of applications per year: 3 (at 10 fl oz/acre of LUNA EXPERIENCE)
- Minimum retreatment interval: 7 days
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- **DO NOT** apply LUNA EXPERIENCE within 35 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- **DO NOT** cut cover crops in treated areas for feed or allow livestock to graze treated areas.

<sup>1</sup>Not for use in CA without a supplemental label.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross-contamination with other pesticides.

**Pesticide Disposal:** Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

### **Container Handling:**

#### **Non-Seed Treatment Products in Non-Refillable Containers**

Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

## **IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE SHALL NOT BE LIABLE FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

LUNA EXPERIENCE is specially formulated and sold by Bayer CropScience LP for the control of various pathogens according to the directions on this label. The purchase price of LUNA EXPERIENCE includes a prepaid license under which purchaser agrees to employ the purchased quantity of LUNA EXPERIENCE only for the above-specified uses and to provide notice of the terms and conditions of this license to any subsequent purchaser. Uses of LUNA EXPERIENCE other than those specified on this label are not licensed through the purchase of this product.

Bayer, Bayer Cross and Luna Experience® are registered trademarks of Bayer Group.







NET CONTENTS: 1/2 Gal. (64 fl. oz.)

# LUNA EXPERIENCE®

FLUOPYRAM	GROUP 7	FUNGICIDE
TEBUCONAZOLE	GROUP 3	FUNGICIDE

Broad spectrum fungicide for control of plant diseases.

Almond; Brassica leafy greens (subgroup 5B); Bulb vegetables (group 3-07); Cucurbits (group 9); Grapes; Hops; Dry and Succulent Beans; Okra; Pistachio; Stone Fruits (group 12); Sunflower; Tree Nuts (group 14).

ACTIVE INGREDIENTS: FLUOPYRAM*	17.6%
TEBUCONAZOLE*	17.6%
OTHER INGREDIENTS:	64.8%
TOTAL:	100.0%

Contains 1.67 lbs FLUOPYRAM and 1.67 lbs TEBUCONAZOLE per gallon

(CAS Numbers 658066-35-4 and 107534-96-3)

EPA Reg. No. 264-1091

SUSPENSION CONCENTRATE

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See additional precautionary statements and directions for use on label.

## PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Produced for:

Bayer CropScience LP

800 N. Lindbergh Blvd.

St. Louis, MO 63167

1-866-99BAYER (1-866-992-2937)

©2022 Bayer CropScience

US86247194D

210915D

07/22



# LUNA EXPERIENCE<sup>®</sup>

## FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

**For MEDICAL Emergencies Call 24 Hours  
A Day 1-800-334-7577.**

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

**NOTE TO PHYSICIAN:** Treat Symptomatically.

**For PRODUCT USE Information Call 1-866-99BAYER  
(1-866-992-2937)**

## DIRECTIONS FOR USE

### STOP - READ THE LABEL BEFORE USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container and keep tightly closed, when not in use. Store in a cool dry place. Avoid cross-contamination with other pesticides.

**Pesticide Disposal:** Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

**Container Handling:** Non-Seed Treatment Products in Non-Refillable Containers

Rigid, Non-refillable containers (equal to or less than 5 gallons) Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill.

**Bayer**