



FLUOPYRAM	GROUP	7	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF CALIFORNIA

LUNA SENSATION®

EPA Reg. No. 264-1090

For the Control of Grapevine Trunk Diseases (GTDs) Through Treating Pruning Wounds in Grape

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Fruit, small vine climbing, except fuzzy kiwifruit, subgroup 13-07F

Amur River Grape; Gooseberry; Grape; Kiwifruit, Hardy; Maypop; Schisandra Berry; cultivars varieties, and/or hybrids of these.

Note: ***DO NOT** apply or allow drift to **Concord grapes** or crop injury may occur. Rinse spray equipment before application of other products to **Concord grapes** or crop injury may occur.*

Disease Control	Application Rate	Application Instructions
Grapevine Trunk Diseases (GTDs) Eutypa dieback (<i>Eutypa lata</i>) Botryosphaeria dieback (<i>Neofusicoccum parvum</i>)	5.0 to 7.0 fl oz/acre (0.082 to 0.115 lb/acre fluopyram) (0.082 to 0.115 lb/acre trifloxystrobin)	Apply within 24 hours of pruning using ground or aerial equipment. Refer to University and/or extension guidelines for best practices and application timings. Continue as needed on a 14- to 21-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Restrictions:

- Maximum single application rate: 7.0 fl oz/acre of LUNA SENSATION (0.115 lb/acre fluopyram and 0.115 lb/acre trifloxystrobin).
- Maximum annual application rate: 21 fl oz of LUNA SENSATION per acre (0.345 lb/acre fluopyram and 0.345 lb/acre trifloxystrobin) per year.
- Maximum number of applications per year: 3 (at 7.0 fl oz/acre of LUNA SENSATION).
- **DO NOT** apply more than 0.446 lbs fluopyram or 0.375 lbs trifloxystrobin per acre per year, including soil and foliar uses, regardless of formulation or method of application.
- **DO NOT** apply LUNA SENSATION after BBCH 73 stage (when bunches begin to hang) or within 45 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, **DO NOT** make more than 2 sequential applications of LUNA SENSATION or any Group 7 or Group 11 containing fungicide before rotating with a fungicide from a different Group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP, 800 N. Lindbergh Blvd., St. Louis, MO 63167

LUNA SENSATION is a registered trademark of Bayer

12/15/2025

Page 1 of 1