













POLLINATOR ADVISORY STATEMENT

This product may adversely impact the forage and habitat of local pollinators, including the monarch butterfly (and its larvae), birds, or bats if reaches non-target areas. Protect pollinators by following label directions to minimize spray drift.

HERBICIDE RESISTANCE MANAGEMENT (WSSA) RECOMMENDATIONS

For resistance management, OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

USE RATES

Winter Wheat and Fall Planted Triticale

Preplant / Pre-emergence Burndown: Apply 0.6 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre with glyphosate in the fall. This product can be mixed with glyphosate containing products in a burn down application in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Follow the glyphosate product label for rate and adjuvant recommendation for control of emerged species.

Postemergence: Apply 0.6 - 0.9 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre with 0.25% - 0.5% non-ionic surfactant in fall or spring in a single application to actively growing weeds. Select rate according to table “Weeds Controlled.”

A fall application of OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE may be followed by a spring application of the product up to a maximum of 1.2 oz per acre OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per year. Apply OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE before jointing begins to avoid crop injury.

Table 1 – Preplant / Pre-emergence Burndown Application with Glyphosate in Winter Wheat for Control or suppression of Bromus species

Timing of Application	OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE Rate* (oz/A)	Remarks
Fall	0.6	Apply 0.6 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre tankmixed with glyphosate as directed for burndown application as a preplant / pre-emergence broadcast spray
Fall followed by Spring	0.6 oz followed by 0.6	Apply 0.6 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre in the fall tankmixed with glyphosate as a preplant burndown followed by 0.6 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre in the spring as a postemergence broadcast spray.

Table 2 – Fall Application with Glyphosate – Planting to spring wheat in following spring

Timing of Application	OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE Rate* (oz/A)	Remarks
Fall	0.4	Apply 0.4 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre tankmixed with glyphosate as directed for burndown application as a preplant / pre-emergence broadcast spray

Table 3 – Postemergence Application for Control or suppression of Bromus species and other weed species shown in “Weeds Controlled”

Timing of Application	Rate* (oz/A)	Remarks
Fall or Spring	0.6 to 0.9	Apply specified dosage per acre as a postemergence broadcast spray
Fall followed by Spring	0.6 – 0.9 oz followed by 0.3 - 0.6 oz	Apply 0.6 – 0.9 oz/A in the fall followed by 0.3 - 0.6 oz/A in the spring as postemergence broadcast sprays. Sequential applications are required for suppression of jointed goatgrass.

*0.6 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre is equal to 0.0263 lbs propoxycarbazone active ingredient per acre. Do not exceed 1.2 oz of OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre or 0.0525 lbs propoxycarbazone active ingredient per acre per year. This limit includes any products, which also contain propoxycarbazone used on the same acre.


Table 4 – Control of wild oat in Idaho, Montana, Oregon, and Washington

Timing of Application	Rate (oz/A)	Remarks
Fall	0.9	Fall emerging wild oat: Use as a fall application before wild oat reaches 2 tillers. Fall applications may not control spring germinating wild oat.
Spring	0.9	Spring emerging wild oat: Apply after emergence but before wild oat reaches the 2-tiller stage of growth.

When used according to label directions, wild oat will be controlled in Idaho, Montana, Oregon, and Washington. Wild oat will be suppressed in areas where OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE is labeled for use outside these four states.

Spring Wheat, Durum Wheat, and Spring Planted Triticale

OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE can be an effective tool in the management of Bromus species in spring wheat and triticale.



Preplant / Pre-emergence Burndown: Apply 0.2 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre with glyphosate as directed for burndown use in fall or spring. Follow the glyphosate product label for rate and adjuvant recommendation for the control of emerged species. **Do not apply preplant or pre-emergence to durum wheat.**

Table 5 – Preplant / Pre-emergence Burndown Application with Glyphosate in Spring Wheat for Suppression of Bromus species		
Timing of Application	OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE Rate* (oz/A)	Remarks
Fall or Early Spring	0.2	Apply 0.2 oz per acre as a preplant / pre-emergence broadcast spray with glyphosate
Fall or Early Spring followed by a Postemergence herbicide, application registered for this use.	0.2 oz followed by a postemergence herbicide treatment registered for this use.	Apply 0.2 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre with glyphosate in the fall or early spring as a preplant burndown. This may be followed as needed with a postemergence herbicide. Please allow a minimum of 15 days between applications.

*0.2 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre is equal to 0.0088 lbs propoxycarbazone active ingredient per acre. A cumulative total of 0.6 oz per acre or 0.0263 lb propoxycarbazone active ingredient per acre may be used per year for spring wheat. This limit includes any products, which also contain propoxycarbazone used on the same acre.

Postemergence in all states except WA, ID and OR: In spring wheat, apply 0.2 - 0.4 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE postemergence per acre in a single application to actively growing weeds. Do not exceed a product application rate of 0.6 oz per acre or 0.0263 lb propoxycarbazone active ingredient per acre per year for spring wheat.

Apply OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE before jointing begins to avoid crop injury.

Postemergence in WA, ID, and OR: In spring wheat, apply OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE at 0.6 – 0.9 oz per acre in a single application to actively growing weeds. Select rate according to table “Weeds controlled.” Do not exceed a product application rate of 0.9 oz per acre or 0.039 lb propoxycarbazone active ingredient per year for spring wheat. Avoid use on ALS-sensitive varieties. Apply OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE before jointing begins to avoid crop injury.

Triticale

Apply 0.6 - 0.9 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre in a single postemergence application to actively growing weeds. **DO NOT APPLY** preplant or PRE Triticale.

See the “**CROP ROTATION RESTRICTIONS**” section for recropping guidance. The crop rotational options will depend on the total amount of propoxycarbazone applied cumulatively in preplant / pre-emergence and/or postemergence treatments.

ENDANGERED SPECIES

To avoid adverse effects on endangered plant species, the following mitigation measures will be required where endangered species occur in Counties listed in the table on the following page.

For ground applications, the applicator must:

1. Apply when there is sustained wind away from native plant communities, OR
2. Leave 50 foot untreated buffer between treatment area and native plant communities.

For aerial applications, the applicator must:

1. Apply only when there is sustained wind away from native plant communities, OR
2. Leave 350 foot untreated buffer between treatment area and native plants.

State	County	State	County	State	County	State	County
Colorado	Morgan	Kansas	Allen	Missouri	Barton	Oregon	Baker
	Weld		Anderson		Benton		Benton
Illinois	Alexander		Bourbon		Cass		Clackamas
	Brown		Coffey		Cedar		Douglas
	Bureau		Crawford		Dade		Jackson
	Calhoun		Douglas		Harrison		Josephine
	Cass		Franklin		Howell		Klamath
	Cook		Jackson		Pettis		Lane
	Du Page		Jefferson		Polk		Linn
	Ford		Johnson		St Clair		Marion
	Fulton		Leavenworth		Vernon		Polk
	Greene		Linn	Nebraska	Cherry		Union
	Grundy		Lyon		Hall		Washington
	Jackson		Miami		Kimball		Yamhill
	Jersey		Neosho		Lancaster	Tennessee	Coffee
	La Salle		Osage	Seward	Dickson		
	Lee		Pottawatomie	New Mexico	Chaves		Lawrence
	Madison	Riley	Ohio		Lucas		Marion
	Marshall	Shawnee		Ottawa	Maury		
	Mason	Kentucky	Barren	Wayne	Williamson		
	Massac		Edmonson	Oklahoma	Craig		Wilson
	Monroe		Fleming		Rogers	Texas	Hidalgo
	Morgan		Grayson		Nueces		
	Ogle		Hardin		Pecos		
	Peoria		Hart	Utah	Emery		
	Pike		Nicholas		Washington	Chelan	
	Putnam					Cowlitz	
	Randolph					Lewis	
	Saline						
	Schuyler						
	Scott						
	St Clair						
	Tazewell						
	Union						
	Winnebago						
	Woodford						

242

234

3

3

5

150

144

3

WEEDS CONTROLLED

OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE effectively controls the following weeds when applied at the rates and application timings shown and weeds are actively growing. Best control is achieved when grass weeds are treated at the 2-leaf to 2-tiller stage of growth and before broadleaf weeds are larger than 2 inches in diameter.

Common name	Scientific name	Fall Application Rates		Spring Application Rates	
		0.6 oz/A	0.9 oz/A	0.6 oz/A	0.9 oz/A
Grasses					
Bulbous bluegrass	<i>Poa bulbosa</i>	S	S	S	S
Cheat (true cheat) ²	<i>Bromus secalinus</i>	C	C	C	C
Dense silky-bent (Windgrass)	<i>Apera spica-venti</i>	C	C	C	C
Downy brome ³	<i>Bromus tectorum</i>	S	C	S	S
Foxtail barley	<i>Hordeum jubatum</i>	C	C	C	C
Hood canarygrass	<i>Phalaris paradoxa</i>	C	C	C	C
Japanese brome ⁴	<i>Bromus japonicus</i>	C	C	C	C
Jointed goatgrass ⁵	<i>Aegilops cylindrica</i>	-	S	-	S
Littleseed canarygrass	<i>Phalaris minor</i>	C	C	C	C
Quackgrass	<i>Elytrigia repens</i>	S	S	S	S
Rattail fescue	<i>Vulpia myuros</i>	S	S	S	S
Rescuegrass	<i>Bromus catharticus</i>	-	S	-	S
Ripgut brome	<i>Bromus rigidus</i>	S	C	S	C
Soft chess	<i>Bromus commutatus</i>	C	C	C	C
Wild oat ⁴	<i>Avena fatua</i>	S	C	S	C
Windgrass	<i>Apera interrupta</i>	C	C	C	C
Broadleaves					
Black mustard	<i>Brassica nigra</i>	C	C	C	C
Blue mustard	<i>Chorispora tenella</i>	C	C	S	C
Burr buttercup	<i>Ranunculus testiculatus</i>	C	C	C	C
Bushy wallflower	<i>Erysimum repandum</i>	S	C	S	C
Field pennycress	<i>Thlaspi arvense</i>	C	C	C	C
Flixweed	<i>Descurania sophia</i>	C	C	C	C
Henbit	<i>Lamium amplexicaule</i>	S	S	-	S
Mouseear chickweed	<i>Cerastium vulgatum</i>	S	C	-	-
Pigweed, redroot	<i>Amaranthus retroflexus</i>	-	S	C	C
Rape (volunteer)	<i>Brassica rapa</i>	C	C	C	C
Shepherdspurse	<i>Capsella bursa-pastoris</i>	C	C	C	C
Small seeded falseflax	<i>Camelina micropora</i>	C	C	C	C
Tall wormseed wallflower	<i>Erysimum cheiranthoides</i>	C	C	C	C
Tansy mustard	<i>Descurania pinnata</i>	C	C	C	C
Tumble mustard	<i>Sisymbrium altissimum</i>	C	C	C	C
Wild buckwheat	<i>Polygonum convolvulus</i>	-	-	-	S
Wild mustard	<i>Brassica kaber</i>	C	C	C	C
Wild turnip	<i>Brassica campestris</i>	C	C	C	C
Note: C means Control S means Suppression					
Suppressed weeds will be stunted in growth and/or be reduced in number as compared to non-treated areas but performance will not be commercially acceptable.					

^{1,2} For field infestations consisting of Japanese brome or cheat (true cheat) only, control may be achieved up to a maximum growth stage of 6-tillers.

³ When weeds are exposed to unfavorable environmental conditions (severe weather conditions, drought, extreme temperatures, etc.), control may be reduced. Applications should be made to actively growing weeds.

⁴ When used according to label directions, wild oat will be controlled in Idaho, Montana, Oregon, and Washington. Wild oat will be suppressed in areas where OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE is labeled for use outside these four states.

⁵ Fall and Spring sequential applications required. See Table 1 application information.

TANKMIXES

For broad-spectrum control of both annual grasses and broadleaf weeds, OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE may be mixed with broadleaf herbicides. With all tank-mix partners, read and follow use directions, rates, precautions, timing and growth stage limitations, recropping restrictions, grazing interval restrictions and recommendations on the broadleaf herbicide and surfactant labels. A non-ionic surfactant is always required with OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE (see “**SURFACTANT**” section).

Possible tank-mix partners include:

Amber®	BRONATE ADVANCED™***	HUSKIE®	Rave™
Aim™	BUCTRIL®***	Harmony® GT XP,	Metribuzin*
Ally XP®, Ally Extra®	Clarity®	Harmony® Extra	Starane™, Starane Flex*,
Affinity Tankmix, Affinity	Curtail®, Curtail® M	MCP Amine or Ester	Starane Ultra™, Starane NXT™
Broadspec®	Finesse®	Orion®	2,4-D Amine or Ester **
Banvel® **		Peak®	WideMatch®

* Spring application in tank-mix combination with Metribuzin may result in reduced control of wild oat.

** Applications with Banvel® or Clarity® (dicamba), Curtail® or 2,4-D Amine or Ester may result in reduced downy brome (*Bromus tectorum*) control.

*** Equivalent bromoxynil products may be substituted

Do not apply OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE in tankmixture with malathion, mancozeb, phosphorodithioate (Di-Syston®), chlorpyrifos (Lorsban®), or methyl parathion as unacceptable crop response may occur.

MIXING INSTRUCTIONS

Ensure the spray tank is clean. In-line strainers and nozzle screens should be clean and 50 mesh or coarser.

1. Fill the spray tank 1/4 to 1/2 full with clean water and begin agitation or bypass.

2. Add the appropriate rate of OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE, as determined under “Recommended Rates”, directly to the spray tank. Maintain sufficient agitation during both mixing and application.

3. Add the broadleaf weed herbicide.

4. Add the surfactant.

5. Fill the spray tank with balance of water needed.

6. Maintain sufficient agitation during both mixing and application of OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE.


RE-SUSPENDING WG PRODUCTS IN SPRAY SOLUTION

OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE may settle if left standing without agitation. If the spray solution is allowed to stand for one hour or more, re-agitate the spray solution for a minimum of 10 minutes before application.

COMPATIBILITY

If OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE is to be tank mixed with other herbicides, compatibility should be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1qt) of spray solution, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually occur within 5-15 minutes after mixing. Read and follow the label of each tank mix product used for precautionary statements, directions for use, geographic and other restrictions. Indications of incompatibility include separation in the mix, and either clumping or clabbering of the mixture.

5



TANK CLEANUP PROCEDURE

1. Drain the tank completely, and then wash out tank, boom and hoses with clean water. Drain again.
2. Half fill the tank with clean water and add ammonia (i.e., 3% domestic ammonia solution) at a dilution rate of 1% (i.e., 1 gallon of domestic ammonia for every 100 gallons of rinsate). Complete filling of the tank with water. Agitate/recirculate and flush through boom and hoses. Leave agitation on for 10 minutes. Drain tank completely.
3. Repeat step 2.
4. Remove nozzles and screens and soak them in a 1% ammonia solution. Inspect nozzles and screens and remove visible residues.
5. Flush tank, boom, and hoses with clean water.
6. Inspect tank for visible residues. If present, repeat step 2.

RE-CROPPING GUIDELINES

OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE breakdown in the soil is due mainly to microbial activity. It can be affected by soil temperature and moisture. Conditions that accelerate the breakdown of OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE include adequate soil moisture and adequate soil temperatures to support microbial activity. Likewise, OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE breakdown can be slowed under dry, cold conditions. When considering crop rotations, soil moisture and soil temperature conditions since application should be monitored.

In determining the appropriate recropping interval to follow when multiple herbicides have been used during the season, follow the label with the longest interval for the crop to be planted from all the herbicides used the previous year.

Recropping intervals following a cumulative total of 0.2 – 0.6 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre

or 0.0088 – 0.0263 lbs propoxycarbazone active ingredient per acre.

The following crops may be recropped after OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE is applied alone or in combination with other products which contain propoxycarbazone.

To ensure safety of rotational crops, the following rotational intervals must be followed:

Minnesota, Montana, North Dakota, South Dakota, Washington, Oregon, and Idaho

Crop	Rotation Interval (Months)
Spring and Winter Wheat	0
Triticale	0
Durum Wheat	4
Proso Millet	4
Grain Sorghum	6
Barley	10
Canola	10
Chickpeas	10
Dry Beans	10
Lentils	10
Peas	10
Soybean	10
Sugarbeets	10
Sunflowers	10
Corn – Conventional	10

Recropping intervals following a cumulative total of 0.6 – 1.2 oz OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE per acre

or 0.0263 – 0.0525 lbs propoxycarbazone active ingredient per acre.

To ensure safety of rotational crops, the following cumulative precipitation and rotational intervals must be followed:

Oklahoma, Kansas, Nebraska, Texas

Crop	Cumulative Precipitation (Inches)	Rotation Interval (Months)
Winter Wheat	0	0
Triticale	0	0
Proso Millet	10	4
Soybean STS™	10	4
Grain Sorghum	10	6
Cotton	24	12
Sunflower	24	12
Soybean - Conventional	24	12
Corn – Conventional	30	18

Washington, Oregon, Idaho, Montana

Crop	Cumulative Precipitation (Inches)	Rotation Interval (Months)
Winter Wheat	0	0
Triticale	0	0
Proso Millet	10	4
Chickpea	24	12
Field Peas	24	12
Spring Barley	24	18
Lentils	24	18
Canola	24	22
Potato	24	22
Corn – Conventional	24	22

Colorado, Minnesota, Wyoming, North Dakota, South Dakota


Crop	Cumulative Precipitation (Inches)	Rotation Interval (Months)
Winter Wheat	0	0
Triticale	0	0
Proso Millet	10	4
Grain Sorghum	10	6
Corn – Conventional	24	22

NOTE: In areas where a crop is not specified or the accumulated precipitation was less than specified above, conduct a field bioassay as described in the “**FIELD BIOASSAY**” section of the label. In all areas, 24-inch rainfall and 24-month rotation interval are required for buckwheat, onions, oats, and sugarbeets.

FIELD BIOASSAY

A field bioassay must be conducted for crops not listed on this label and for crops where cumulative precipitation requirements are not satisfied or for crops listed on the label for which a shorter plant-back interval than listed is desired. To conduct a field bioassay, plant strips of the crop you want to grow the season following OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE application. Monitor the crop for response to OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE to determine if the crop can be grown safely in previously treated OLYMPUS 70% WATER DISPERSIBLE GRANULAR HERBICIDE areas.

Regardless of the bioassay results, do not plant any crop, except wheat, closer than 4 months after application.



242

234

5

3

3

150

144

3

OLYMPUS® 70% Water Dispersible Granular Herbicide

For control of certain grasses and broadleaf weeds in wheat and triticale.

ACTIVE INGREDIENT: Propoxycarbazone-sodium

OTHER INGREDIENTS

TOTAL:

70.00%

30.00%

100.00%

EPA Reg. No. 264-809

EPA Est. 000264-DEU-001

STOP - Read the label before use

KEEP OUT OF REACH OF CHILDREN

CAUTION PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail)

See Back Panel for First Aid Instructions and Booklet for Complete Precautionary Statements and Directions for Use.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF SWALLOWED:	- Call a poison control center or doctor immediately for treatment advice. - Do not induce vomiting unless told to do so by a poison control center or doctor. - Have person sip a glass of water if able to swallow. - Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.
IF INHALED:	- Move person to fresh air. - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth to mouth. - Call a poison control center or doctor for further treatment.
IF IN EYES:	- Hold eye open and rinse slowly and gently with water for 15-20 minutes. - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. - Call a poison control center or doctor for treatment advice.

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

Drift or runoff may adversely affect non-target plants.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer CropScience Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer CropScience Emergency Response Telephone No. is 1-800-334-7577.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Bayer and Olympus are registered trademarks of Bayer Group.

©2026 Bayer Group. All rights reserved.

Produced for:
Bayer CropScience LLC
800 N. Lindbergh Blvd.
St. Louis, MO 63167

Bayer

Net Contents: 1 lb (454 grams)

US89141524E 251120Ev2 08/26