



FIFRA Section 24(c) Special Local Need (SLN) Label
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

PROLINE[®] 480 SC Fungicide

For Suppression of Black Leg (*Leptosphaeria maculans*) and Chlorotic leaf spot (*Cylindrosporium concentricum*) in Brassicaceae Grown For Seed

EPA Reg. No. 264-825

EPA SLN. No. OR-150017

THIS LABEL IS VALID UNTIL DECEMBER 31, 2030, UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

ACTIVE INGREDIENT:

Prothioconazole: 2-[2-(1-Chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-1, 2-dihydro-3H-1,2,4-triazole-3-thione 41.0%

OTHER INGREDIENTS 59.0%

Contains 4 pounds prothioconazole per gallon. **TOTAL:** 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the label affixed to the main container of PROLINE 480 SC Fungicide must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the label affixed to the main container of PROLINE 480 SC Fungicide and this SLN label.

Brassicaceae Grown For Seed Production, including:

Brassica Grown for Seed (Broccoli; broccoli, Chinese; broccoli raab; Brussels sprouts; cabbage; cabbage, Chinese, bok choy; cabbage, Chinese, napa; cauliflower; collards; kale; rape greens; rutabaga; turnip)

Raphanus (daikon or radish) Grown for Seed

Sinapis (white and yellow mustard) Grown for Seed

Eruca (arugula) Grown for Seed

PEST SUPPRESSION	RATE OF PROLINE 480 SC FUNGICIDE
Black Leg (<i>Leptosphaeria maculans</i>) Chlorotic leaf spot (<i>Cylindrosporium concentricum</i>)	5.0 – 5.7 fl oz/A (0.156 – 0.178 lb ai/A)

RECOMMENDED APPLICATION TIMINGS FOR BIENNIAL BRASSICACEAE SEED CROPS:

Up to 2 applications in the fall may be followed with up to 3 applications in the spring/summer, using the 5.0 fl oz/acre (0.156 lb ai/A) rate.

- **Fall Applications:** Apply PROLINE 480 SC Fungicide in the fall just prior to the onset of cool and wet conditions that favor spread of black leg, particularly if the disease has been detected in the crop.
- **Spring Applications:** Apply PROLINE 480 SC Fungicide in the spring when conditions become warm enough for the plants to start growing again and are moist and cool (particularly west of the Cascade Mountains) – 1 or 2 applications may be needed depending on disease pressure and environmental conditions.
- **Summer Applications:** Apply PROLINE 480 SC Fungicide in the summer when pods begin to form and conditions are cool and moist, particularly if black leg has been detected in the crop earlier in the season and there is a risk of infection occurring on seed developing in the pods.

RESTRICTIONS FOR BIENNIAL BRASSICACEAE SEED CROPS:

- **DO NOT** make more than 5 applications of PROLINE 480 SC Fungicide per acre per year.
- **DO NOT** apply more than 25.0 fl oz/acre (0.781 lb ai/A) of PROLINE 480 SC Fungicide per acre per year.

RECOMMENDED APPLICATION TIMINGS FOR ANNUAL BRASSICACEAE SEED CROPS:

Make up to 3 applications in the spring/summer, using the 5.7 fl oz/acre (0.178 lb ai/A) rate.

- **Spring Applications:** Apply PROLINE 480 SC Fungicide after plants have emerged and conditions are moist and cool (particularly west of the Cascade Mountains) – 1 or 2 applications may be needed depending on disease pressure and environmental conditions.
- **Summer Applications:** Apply PROLINE 480 SC Fungicide in the summer when pods begin to form and conditions are cool and moist, particularly if black leg has been detected in the crop earlier in the season and there is a risk of infection occurring on seed developing in the pods.

RESTRICTIONS FOR ANNUAL BRASSICACEAE SEED CROPS:

- **DO NOT** make more than 3 applications of PROLINE 480 SC Fungicide per acre per year.
- **DO NOT** apply more than 17.1 fl oz/acre (0.534 lb ai/A) of PROLINE 480 SC Fungicide per acre per year.

RESTRICTIONS

- PROLINE 480 SC Fungicide may be applied by ground or air. Apply in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply to Brassicaceae grown for food or feed, nor allow plant components to be utilized for human food or animal feed consumption.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** windrow within 14 days of last application.

NOTES

Use in rotation in resistance management programs where additional products are available. Please refer to the PROLINE 480 SC Fungicide Section 3 label for additional information

Special Crop Use Restrictions

The pesticide applicator, the producer of these crops, and the seed conditioner must be aware that use of this product according to this labeling is deemed a non-feed/non-food use by the Oregon Department of Agriculture, and is regulated by the Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of these crops who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for these crops. Consequently, no portion of these seed crops may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to, green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies."

IMPORTANT—READ BEFORE USE

PROLINE 480 SC FUNGICIDE, WHEN USED ON BRASSICACEAE GROWN FOR SEED UNDER THIS SECTION 24(C) LABEL HAS NOT BEEN TESTED UNDER ALL CONDITIONS AND IN ALL TANKMIX SCENARIOS. BAYER RECOMMENDS THAT THE USER AND/OR GROWER TEST PROLINE 480 SC FUNGICIDE IN ORDER TO DETERMINE ITS COMPATIBILITY WITH THE INTENDED USE. **BECAUSE OF THE RISK OF CROP INJURY, CROP YIELD REDUCTION, AND/OR CROP LOSS, ALL SUCH USE IS AT USER'S AND/OR GROWER'S RISK.**

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For **PRODUCT USE** Information Call **1-866-99BAYER (1-866-992-2937)**.

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

24(c) Registrant:

**Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167**