



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

PROLINE® 480 SC Fungicide

EPA Reg. No.: 264-825

For use on Corn

Supplemental Label

PROLINE® 480 SC Fungicide

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product package label for **PROLINE® 480 SC Fungicide**.

USE DIRECTIONS

RECOMMENDED APPLICATIONS		
CROP	DISEASE CONTROLLED	RATE OF PROLINE 480 SC FUNGICIDE
CORN (Field Corn, Field Corn Grown for Seed, Sweet Corn and Popcorn)	Anthrachnose Leaf Blight (<i>Colletotrichum graminicola</i>) Eye Spot (<i>Aureobasidium zeae</i>) Gray Leaf Spot (<i>Cercospora zeae-maydis</i>) Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>)* Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>)* Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>)* Rusts (<i>Puccinia</i> spp.) *The above diseases are also known as Helminthosporium leaf blights	5.7 fl oz. per acre
<p>General Comments: PROLINE® 480 SC may be applied by either ground, aerial or chemigation application equipment. For aerial applications, PROLINE 480 SC may be applied using a minimum of 2 gpa spray solution.</p> <p>Apply PROLINE 480 SC at the first sign of disease. Repeat applications as needed on a 7- to 14-day interval if favorable conditions for disease development persist. Application of PROLINE 480 SC is not recommended at times when corn is under severe environmental stress conditions.</p> <p>Do not apply more than 22.8 fl. oz. of PROLINE 480 SC per acre per crop. For field corn, field corn grown for seed and popcorn, do not apply within 14 days of harvest for grain and fodder. Forage may be harvested the same day of application. For sweet corn, do not apply within 14 days of harvest for fodder. Sweet corn ears and forage may be harvested the same day of application.</p>		

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).