



FLUOPYRAM	GROUP	7	FUNGICIDE
PROTHIOCONAZOLE	GROUP	3	FUNGICIDE

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE OF STATES OF ALABAMA, ARKANSAS, FLORIDA, GEORGIA, MISSISSIPPI, NORTH CAROLINA, NEW MEXICO, OKLAHOMA, SOUTH CAROLINA, TEXAS AND VIRGINIA

This recommendation is valid until December 31, 2028

PROPULSE®

EPA Reg. No. 264-1084

To allow for in furrow application to peanuts for the control of Early Leaf Spot, Late Leaf Spot, and Nematodes.

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

PEANUT

Disease/Pest Control	Application Rate	Application Instructions
Early Leaf Spot (<i>Cercospora arachidicola</i>) Late Leaf Spot (<i>Cercosporidium personatum</i>) Nematodes	13.6 fl oz/acre (0.177 lb/acre fluopyram) (0.177 lb/acre prothioconazole)	May be applied in furrow

Restrictions:

- Maximum single application rate: 13.6 fl oz/acre of PROPULSE (0.177 lb/acre fluopyram and 0.177 lb/acre prothioconazole)
- Maximum annual application rate: 27.2 fl oz/acre of PROPULSE (0.354 lb/acre fluopyram and 0.354 lb/acre prothioconazole).
- Maximum number of applications per year: 2
- Minimum retreatment interval: 14 days
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs fluopyram or 0.712 lbs prothioconazole per acre per year from all uses, including seed treatment, soil and foliar applications.
- Apply by ground, aerial, or chemigation application equipment.
- Minimum application volumes: 10 **gallons/Acre** (Ground); 5 **gallons/Acre** (Aerial).
- Pre-Harvest Interval (PHI): 14 days
- **DO NOT** feed hay or threshings or allow livestock to graze in treated areas.
- To limit the potential for development of disease resistance to this fungicide, **DO NOT** make more than 2 sequential applications of PROPULSE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- **DO NOT** use with handheld application equipment, including mechanically pressurized spray gun, backpack or tank pressurized spray gun, or handheld boom applicators.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP, 800 N. Lindbergh Blvd., St. Louis, MO 63167

Propulse is a registered trademark of Bayer

1/27/2026

Page 1 of 1