



FLUOPYRAM	GROUP	7	FUNGICIDE
PROTHIOCONAZOLE	GROUP	3	FUNGICIDE

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ALABAMA, ARKANSAS, COLORADO, DELAWARE, FLORIDA, GEORGIA, IDAHO, KANSAS, MARYLAND, MISSISSIPPI, NEBRASKA, NEW JERSEY, NORTH CAROLINA, NORTH DAKOTA, OKLAHOMA, OREGON, PENNSYLVANIA, SOUTH CAROLINA, TEXAS, VIRGINIA, WASHINGTON, AND WISCONSIN

This recommendation is valid until December 31, 2026.

PROPULSE®

EPA Reg. No. 264-1084

For Suppression of Nematodes at a reduced in-furrow application rate in Corn

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Tank mixes with Propulse and some fertilizers and micronutrients have been problematic. It is highly recommended that users conduct a “jar-test” to make sure that tank mix partners are compatible before mixing. Contact your local Bayer Rep for more information.

CORN Sweet Corn; Field Corn; Field Corn Grown For Seed, and Popcorn		
Pest Suppression	Application Rate	Application Instructions
Nematodes	8.0 fl oz/acre	In-furrow spray during planting directed on or below seed. Tank mixes with Propulse and some fertilizers and micronutrients has been problematic. It is highly recommended that users conduct a “jar-test” to make sure that tank mix partners are compatible before mixing. Contact your local Bayer rep for more information.

CORN - Continued

Restrictions:

- Maximum single application rate: 8.0 fl oz/acre of PROPULSE (0.104 lb/acre fluopyram and 0.177 lb/acre prothioconazole)
- Maximum annual application rate: 27.2 fl oz of PROPULSE per acre (0.354 lb/acre fluopyram and 0.354 lb/acre prothioconazole) per year.
- Maximum number of applications per year: 3
- Minimum retreatment interval: 7 days
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lbs fluopyram or 0.712 lbs prothioconazole per acre per year from all uses including seed treatment, soil and foliar applications.
- Minimum application volumes: **10 gallons/Acre** (Ground)
- Apply by either ground or chemigation application equipment.
- Pre-Harvest Interval (PHI): 0 day - sweet corn ears and forage, field corn forage, and field corn grown for seed; 14 days – grain and fodder of field corn and popcorn.
- For sweet corn, the Restricted-Entry Interval (REI) is 24 hours.
- To limit the potential for development of disease resistance to this fungicide, **DO NOT** make more than 2 sequential applications of PROPULSE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- **DO NOT** use with handheld application equipment, including mechanically pressurized spray gun, backpack or tank pressurized spray gun, or handheld boom applicators.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167