



Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, NC 27709
(919) 549-2000

PROSARO® 421 SC Fungicide
EPA Reg. No. 264-862

Use on Wheat - Reduced Application Rate for
Management of Rust Diseases

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF KANSAS

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee). It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

CROP	DISEASE CONTROLLED	RATE OF PROSARO 421 SC Fungicide fl oz/acre
Wheat	RUSTS (<i>Puccinia spp.</i>)	5.0
Apply PROSARO® 421 SC Fungicide when the earliest disease symptoms appear on the leaves or stems. Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. For optimum disease control, the lowest labeled rate of a spray surfactant must be tank-mixed with PROSARO 421 SC Fungicide. Restrictions: - Do not apply within 30 days of harvest. Straw may be fed or used for bedding. - Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with PROSARO 421 SC Fungicide.		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at <http://www.bayercropscience.us>

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).