



PROTHIOCONAZOLE	GROUP	3	FUNGICIDE
TEBUCONAZOLE	GROUP	3	FUNGICIDE
FLUOPYRAM	GROUP	7	FUNGICIDE

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF CONNECTICUT, KANSAS, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, NEBRASKA, NEW HAMPSHIRE, NEW JERSEY, OHIO, PENNSYLVANIA, RHODE ISLAND, VERMONT, WEST VIRGINIA

PROSARO® PRO 400 SC Fungicide (EPA Reg. No. 264-1209)

+

PROLINE® 480 SC Fungicide (EPA Reg. No. 264-825)

For the Tank Mix Application of PROSARO PRO 400 SC + PROLINE 480 SC Fungicides for Suppression of Fusarium and Gibberella Ear Rot on Field Corn, Field Corn Grown for Seed, and Popcorn

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its label.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this 2(ee) recommendation and the EPA registered label for PROSARO PRO 400 SC Fungicide and PROLINE 480 SC Fungicide.

Apply PROSARO PRO 400 SC + PROLINE 480 SC Fungicides at the following rates to suppress fusarium and Gibberella ear rot on field corn, field corn grown for seed, and popcorn:

Rate of PROSARO PRO 400 SC Fungicide	Rate of PROLINE 480 SC Fungicide
10.3 fl oz/A (0.134 lb ai/A prothioconazole) (0.067 lb ai/A tebuconazole) (0.067 lb ai/A fluopyram)	1.41 fl oz/A (0.044 lb ai/A prothioconazole)

For optimum suppression of fusarium and Gibberella ear rot, apply **PROSARO PRO 400 SC + PROLINE 480 SC Fungicides** from R1 (initial silk emergence) to the R2 (brown silk) corn growth stages. This combination will reduce both disease symptoms and levels of mycotoxin in the grain.

RESTRICTIONS

- Pre-Harvest Interval (PHI) for forage: **21 day(s)**. **DO NOT** allow livestock to graze corn forage within 21 days of application.
- Pre-Harvest Interval (PHI) for grain or fodder: **14 day(s)**. **DO NOT** allow livestock to graze corn fodder within 36 days of application.
- Minimum interval between applications: **14 days**
- Minimum application volumes: **10 gallons/Acre** (Ground); **2 gallons/Acre** (Aerial)

- **DO NOT APPLY** more than 41.2 fluid ounces/Acre (0.538 lbs prothioconazole, 0.270 lbs tebuconazole, 0.270 lbs fluopyram) per acre of PROSARO PRO 400 SC Fungicide allowed per year.
- **DO NOT APPLY** more than 22.8 fl oz (0.713 lbs prothioconazole) per acre of PROLINE 480 SC Fungicide per year.
- Regardless of formulation or method of application, **DO NOT** exceed 0.713 lb prothioconazole, 0.675 lb tebuconazole, or 0.446 lb fluopyram per acre per year from all uses, including seed treatment, soil and foliar applications.
- To limit the potential for development of disease resistance to this fungicide, **DO NOT** make more than 2 sequential applications of PROSARO PRO 400 SC Fungicide or any Group 7 containing fungicide before rotating with a fungicide from a different Group.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

**Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167**