



RIMFIRE[®] MAX

Meet Your Challenging Weed Spectrum Head-On

Rimfire[®] Max Herbicide controls your problem grasses and broadleaf weeds in wheat.

/// BENEFITS

- **Weed Control That Doesn't Miss**
Rimfire Max handles more than 30 tough-to-control grass and broadleaf weeds.
- **Crop Rotation Flexibility**
The rotational flexibility of Rimfire Max is excellent with a 10-month rotation back to 12 key crops in certain geographic regions.
- **Tankmix Partner Flexibility**
Rimfire Max can be mixed with broadleaf herbicides and a non-ionic surfactant.

/// APPLICATION RATES

- Target weeds should be actively growing at time of application.
- Weed control may be reduced when weeds are under stress due to severe weather conditions, drought, very cold temperatures, etc.
- Apply 3 oz. Rimfire[®] Max Herbicide with selected adjuvant per acre to wheat in the spring as a single application.
- Do not use more than 3 oz of Rimfire Max per acre.

/// APPLICATION WINDOW

- Wheat: From emergence - fully expanded first true leaf - up to emergence of flag leaf.
- Grass weeds: 1-leaf up to 2-tiller stage.
- Broadleaf weeds: Treat weeds up to 2 inches in height.
- Do not use more than 3 oz of Rimfire Max per acre.

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

Registered Crops

- | | |
|----------------|----------------|
| • Spring Wheat | • Winter Wheat |
| • Durum | |

Key Weeds Controlled

- | | |
|-------------------|----------------------|
| • Barnyardgrass | • Blue Mustard |
| • Foxtail Barley* | • Black Mustard |
| • Green Foxtail* | • Canola (volunteer) |
| • Japanese Brome* | • London Rocket |
| • Wild Beet | • Tansy Mustard |
| • Wild Oat | • Tumble Mustard |
| • Yellow Foxtail* | • Wild Mustard |

For a complete list of weeds controlled visit [Cereal Experts.com](http://CerealExperts.com)

* Rimfire Max applied at the pre-tiller stage of growth

Application Rate

3 oz/A

Mode-of-Action Group

② Thien carbazone

Recommended Spray Volume

10 - 15 gal/A*

*For aerial application minimum spray volume of 5 gals/A.



LEARN MORE

RIMFIRE[®]
MAX



*For more information on herbicide group numbers and mode-of-action options, visit IWillTakeAction.com

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Bayer, Bayer Cross and Rimfire[®] are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2025 Bayer Group. All rights reserved.