





US79884087G (250315Gv2) RIMFIRE MAX 90 oz. ECL DRUM LABEL

Colors: CMYK Date: 06/16/2026 (Label Coordinator: Louis Cross)

















145 mm

139 mm

130 mm

120 mm

WEEDS CONTROLLED

RIMFIRE Max Herbicide effectively controls the following weeds¹ when applied at the rates and application timings shown and weeds are actively growing. Best control is achieved when grass weeds are treated at the 1-leaf to 2-tiller stage of growth and before broadleaf weeds are larger than 2 inches in height.

GRASS WEED RECOMMENDATIONS		
Cheat (True cheat)*	<i>Bromus secalinus</i>	C
Barnyardgrass*	<i>Echinochloa crus-galli</i>	C
Downy brome	<i>Bromus tectorum</i>	PC
Foxtail barley*	<i>Hordeum jubatum</i>	C
Green foxtail*	<i>Setaria viridis</i>	C
Japanese brome*	<i>Bromus japonicus</i>	C
Little barley	<i>Hordeum pusillum</i>	PC
Persian darnel*	<i>Lolium persicum</i>	PC
Quackgrass	<i>Agropyron repens</i>	PC
Smooth brome	<i>Bromus inermis</i>	PC
Wild oat	<i>Avena fatua</i>	C
Yellow Foxtail*	<i>Setaria pumila</i>	C

BROADLEAF WEED RECOMMENDATIONS		
Blue mustard	<i>Chorispora tenella</i>	C
Black mustard	<i>Brassica nigra</i>	C
Catchweed bedstraw	<i>Galium aparine</i>	PC
Canola (volunteer)	<i>Brassica napus</i>	C
Common chickweed	<i>Stellaria media</i>	PC
Cornflower / Bachelor's Button	<i>Centaurea cyanus</i>	PC
Dogfennel	<i>Eupatorium capillifolium</i>	PC
Field pennycress	<i>Thlaspi arvense</i>	PC
Henbit	<i>Lamium amplexicaule</i>	PC
Ivyleaf speedwell	<i>Veronica hederifolia</i>	PC
London rocket	<i>Sisymbrium irio</i>	C
Mouseear chickweed	<i>Cerastium vulgatum</i>	PC
Red clover	<i>Trifolium pratense</i>	PC
Redroot pigweed ¹	<i>Amaranthus retroflexus</i>	PC
Shepherd's purse	<i>Capsella bursa-pastoris</i>	PC
Swinecress	<i>Coronopus didymus</i>	PC
Tansy mustard	<i>Descurania pinnata</i>	C
Tumble mustard	<i>Sisymbrium altissimum</i>	C
Wild beet	<i>Beta vulgaris</i>	C
Wild mustard	<i>Sinapis arvensis</i>	C
Wild radish	<i>Raphanus raphanistrum</i>	C
Wild turnip	<i>Brassica rapa</i>	C

C means Control. PC means Partial Control. Partially controlled weeds will be stunted in growth and/or be reduced in number as compared to non-treated areas but performance will not be commercially acceptable.

* These weeds will be controlled when RIMFIRE Max Herbicide is applied at the pre-tiller stage of weed growth.


¹ Naturally occurring resistant biotypes of weeds are known to occur. Refer to the **WEED RESISTANCE** section for additional information regarding management tactics for resistant weeds.


USE RATES


Unless otherwise recommended by Bayer CropScience, do not use less than 3 oz per acre of RIMFIRE Max Herbicide. Apply RIMFIRE Max Herbicide at 3 ounces/acre to wheat in spring as a single application to actively growing weeds. Do not exceed a product application rate of 3 ounces/acre in a single application in the spring.

TANKMIXES

For broad-spectrum control of both annual grasses and broadleaf weeds, RIMFIRE Max Herbicide may be mixed with broadleaf herbicides and a non-ionic surfactant. RIMFIRE Max Herbicide contains 0.144 pounds of mefenpyr-diethyl per pound of product. Applying the maximum-labeled rate of RIMFIRE Max Herbicide delivers 0.027 lbs of mefenpyr-diethyl per acre. Do not apply more than 0.053 pounds of mefenpyr-diethyl per acre per year. With all tank-mix partners, read and follow use directions, rates, precautions, timing, recropping restrictions, grazing interval restrictions and recommendations on the broadleaf herbicide and surfactant labels.







To conduct a field bioassay, plant strips of the crop you want to grow the season following RIMFIRE Max Herbicide application. Monitor the crop for response to RIMFIRE Max Herbicide to determine if the crop can be grown safely in previously treated RIMFIRE Max Herbicide areas.

Regardless of the bioassay results, do not plant any crop, except fall-sown or winter wheat, closer than 4 months after a RIMFIRE Max Herbicide application.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications:

- When applying aerially to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- When applying to crops via aerial application equipment, the spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip or rotor blade vortices. The boom length must not exceed 65% of the wingspan of airplanes or 75% of the rotor blade diameter of helicopters.
- When applying to crops via aerial application equipment, applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Ground Boom Applications:

- When using ground application equipment, apply with nozzle height no more than 3 feet above the ground or crop canopy.
- Applicators are required to use a coarse or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.


IMPORTANCE OF DROPLET SIZE


The most effective way to reduce drift potential is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.



Controlling Droplet Size – Ground Boom



- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.











US79884087G (250315Gv2) RIMFIRE MAX 90 oz. Base DRUM LABEL

Colors: Black Date: 06/16/2026 (Label Coordinator: Louis Cross)

