



FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF CA, FL, and TX

This recommendation is valid until December 31, 2028

SIVANTO[®] prime

EPA Reg. No. 264-1141

For the control of Cottony Cushion Scale (foliar) and Glassy-Winged Sharpshooter (foliar) and the suppression of California Red Scale (foliar) and Citrus Leafminer (soil) in Citrus Fruits

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

CITRUS FRUITS (Group 10-10) FOLIAR

Calamondin; Citron; Citrus hybrids; Grapefruit; Japanese summer grapefruit; Kumquat; Lemon; Lime; Lime (Sweet, Australian desert, Australian finger, Australian round, Brown River finger, Mount White, New Guinea wild, Russell River, Tahiti); Mediterranean mandarin; Orange (sour, sweet, Tachibana, Trifoliata); Pummelo; Satsuma mandarin; Tangelo; Tangerine (Mandarin); Tangor; Uniq fruit; cultivars, varieties, and/or hybrids of these

Pest Controlled	Product Rate (fl oz/A)
Cottony cushion scale	14.0
Glassy-winged sharpshooter	10.0 – 14.0
Pest Suppressed	Product Rate (fl oz/A)
California red scale	14.0

Application Restrictions:

Pre-Harvest Interval (PHI): **1 Day(s)**

Minimum interval between applications: **10 Day(s)**

Minimum Application Volume: **50 Gallons/Acre** | Ground; **10 Gallons/Acre** | Aerial

Do not apply more than **28 fl oz** of SIVANTO prime (**0.365 Pound Flupyradifurone**) per **Acre Per Calendar Year** on citrus fruit, regardless of method of application, product or formulation.

CITRUS FRUITS (Group 10-10)
SOIL

Calamondin; Citron; Citrus hybrids; Grapefruit; Japanese summer grapefruit; Kumquat; Lemon; Lime; Lime (Sweet, Australian desert, Australian finger, Australian round, Brown River finger, Mount White, New Guinea wild, Russell River, Tahiti); Mediterranean mandarin; Orange (sour, sweet, Tachibana, Trifoliolate); Pummelo; Satsuma mandarin; Tangelo; Tangerine (Mandarin); Tangor; Uniq fruit; cultivars, varieties, and/or hybrids of these

Pest Suppressed	Product Rate (fl oz/A)
Citrus leafminer	28.0

Application Information:

SIVANTO prime may be applied at the specified dosage by the following methods:

- Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment
- Basal drench in sufficient water to move SIVANTO prime into root zone

Application Restrictions:

Pre-Harvest Interval (PHI): **30 Day(s)**

Do not apply more than **28 fl oz** of SIVANTO prime (**0.365 Pound Flupyradifurone**) per **Acre Per Calendar Year** on citrus fruit, regardless of method of application, product or formulation.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167