



FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ALABAMA, ARIZONA, ARKANSAS, COLORADO, FLORIDA, GEORGIA, KANSAS, KENTUCKY, LOUISIANA, MISSISSIPPI, NEBRASKA, NEW MEXICO, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE, TEXAS, AND VIRGINIA

This recommendation is valid until December 31, 2028

SIVANTO[®] prime

EPA Reg. No. 264-1141

Reduced rates in Pearl Millet and Sorghum for control of Aphids

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

MILLET, PEARL; SORGHUM (INCLUDING SORGHUM GROWN FOR SYRUP) – Foliar Application	
Pest Controlled	Product Rate (fl oz/A)
Aphids	4.0 – 7.0
Application Restrictions: Pre-Harvest Interval (PHI): 7 Day(s) – hay, forage, sorghum grown for syrup, or sweet corn Pre-Harvest Interval (PHI): 21 Day(s) – dried grain, stover or straw Minimum interval between applications: 7 Day(s) Minimum Application Volume: 10 Gallons/Acre Ground; 3 Gallons/Acre Aerial Do not apply more than 28 fl oz of SIVANTO prime (0.365 Pound Flupyradifurone) per Acre Per Calendar Year on cereal grains, regardless of method of application, product or formulation.	

SORGHUM (INCLUDING SORGHUM GROWN FOR SYRUP) – Soil Application

Pest Controlled	Product Rate (fl oz/A)
Aphids	6.0 – 10.0

Application Information:

SIVANTO prime may be applied at the specified dosage by the following methods:

- Injection below the eventual seed row prior to planting. Place SIVANTO prime 3 - 4 inches below seed-line
- In-furrow spray during planting
- Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment

Application Restrictions:

Pre-Harvest Interval (PHI): **45 Day(s)**

Do not apply more than **28 fl oz** of SIVANTO prime (**0.365 Pound Flupyradifurone**) per **Acre Per Calendar Year** on Sorghum, regardless of method of application, product or formulation.

Maximum number of crop seasons per year: **1**

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).



Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167