



FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ALABAMA, ARKANSAS, FLORIDA, GEORGIA, LOUISIANA, MISSISSIPPI, MISSOURI, NEW MEXICO, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE AND VIRGINIA

VELUM®

EPA Reg. No. 264-1078

For reduced application rate in Peanut

This recommendation is valid until December 31, 2029.

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

PEANUT		
Disease/Pest Suppression	Application Rate	Application Instructions
Early leaf spot (<i>Cercospora arachidicola</i>) Late leaf spot (<i>Cercosporidium personatum</i>)	3.25 fl oz/acre	Soil Application Apply specified dosage using any of the following methods: • In-furrow spray during planting directed on or below seed. • Chemigation into root-zone through low-pressure drip or trickle irrigation. Minimum 14-day interval between soil applications.
Aspergillus crown rot (<i>Aspergillus niger</i>) White mold (<i>Sclerotium rolfsii</i>)	4 fl oz/acre	
Restrictions: • Maximum single application rate: 6.84 fl oz/acre of VELUM (0.222 lb/acre fluopyram) • Maximum annual application rate: 13.7 fl oz of VELUM per acre (0.445 lb/acre fluopyram) per year. • Maximum number of applications per year: 2 (at 6.84 fl oz/acre of VELUM) • Minimum retreatment interval: 14 days • DO NOT apply more than 0.446 lbs Fluopyram per acre per year, regardless of formulation or		

method of application (seed treatment, soil or foliar).

- Apply using ground or chemigation equipment.
- **DO NOT** apply VELUM within 7 days of harvest.
- To limit the potential for development of disease resistance to this fungicide class, **DO NOT** make more than 2 sequential applications of VELUM or any Group 7-containing fungicide before rotating with a fungicide from a different Group.
- For soil application, to limit the potential for development of disease resistance to this chemical class, it is advised that the first foliar fungicide application after VELUM be a product from a different FRAC group.
- **DO NOT** allow livestock to graze peanut hay.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167