



VELUM[®]

RISE

Net Contents:

2.5 Gallons

FLUOPYRAM	GROUP	7	FUNGICIDE
PENFLUFEN	GROUP	7	FUNGICIDE

Fungicide and nematicide for use in potato as a soil treatment

ACTIVE INGREDIENTS:

Fluopyram*: 22.12%

Penflufen*: 9.38%

OTHER INGREDIENTS: 68.50%

TOTAL: 100.00%

Contains 2.08 lbs FLUOPYRAM and 0.88 lbs PENFLUFEN per U.S. gallon

*CAS Numbers 658066-35-4 (fluopyram) and 494793-67-8 (penflufen)

EPA Reg. No. 264-1222

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for First Aid Instructions and Booklet for Complete
Precautionary Statements and Directions for Use.

For MEDICAL and TRANSPORTATION Emergencies ONLY

Call 24 Hours a Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

Produced for
Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167

Velum is a registered trademark of Bayer Group.

©2025 Bayer Group. All rights reserved.

FIRST AID

IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • DO NOT give anything by mouth to an unconscious person.
----------------------	---

IF ON SKIN or CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
--------------------------------	---

In case of emergency, call the toll-free Bayer CropScience Emergency Response telephone number: 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Treat Symptomatically.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, and waterproof or chemical-resistant gloves composed of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of Fluopyram. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

Ground Water Advisory

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOURCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSOURCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

STOP - READ THE LABEL BEFORE USE

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements, specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for sale, distribution, or use in Nassau and Suffolk counties, New York except as permitted under FIFRA 24(c), Special Local Need registration.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the **restricted-entry interval (REI)** of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over long-sleeved shirt and long pants, socks and shoes, and chemical-resistant gloves composed of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils.

PRODUCT INFORMATION

VELUM RISE is a broad-spectrum fungicide and nematocide for use in potatoes as a soil treatment for control or suppression of listed crop diseases and control or suppression of plant pathogenic nematodes.

RESISTANCE MANAGEMENT

For resistance management, please note that VELUM RISE contains a Group 7 fungicide. Any fungal population may contain individuals naturally resistant to VELUM RISE and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these (fungicides) are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of VELUM RISE or other Group 7 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

IPM: It is advised that applications of fungicides be integrated into an overall disease and pest management program, following cultural practices known to reduce disease development. Consult your local extension specialist, certified crop advisor and/or manufacturer representative for additional IPM strategies established for your area. This product may be used in Agricultural Extension advisory (disease forecasting or risk assessment) programs, which specify application timing based on environmental factors favorable for disease development.

MANDATORY SPRAY DRIFT

Ground Applications

- For all applications, applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.
- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.

Boom-less Ground Applications

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversion.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

BOOM-LESS GROUND APPLICATIONS

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

HANDHELD TECHNOLOGY APPLICATIONS

Take precautions to minimize spray drift.

APPLICATION INFORMATION

In-furrow at-plant applications

Apply in-furrow during planting operations. Direct applications into the open furrow and cover with soil.

Pre-plant incorporated applications

Apply as a broadcast application prior to planting operations and incorporate into soil using a rototiller or disc or similar implement.

SPRAY MIXING AND COMPATIBILITY

Begin with clean spray equipment and add one-half of the required amount of water to the spray or mixing tank and start agitation. Add the required quantity of fungicide and the tank-mix partner, if applicable to the water and complete filling with water to the required total volume. Follow the advice of your State Cooperative Extension Service for tank mixing with other products. In

general, follow the order beginning first with water conditioners, water soluble packaging (wait for it to completely dissolve), wettable powders and water dispersible granular products, liquid flowables and suspension concentrates, emulsifiable concentrates, and adjuvants last. Maintain agitation throughout spraying. **DO NOT** allow spray mixture to remain in the tank overnight, or for long periods during the day without agitation.

VELUM RISE is physically compatible with most commonly used fungicide, herbicide, insecticide, and foliar nutrient products. However, the compatibility of VELUM RISE with all potential tank-mix partners has not been fully investigated. Compatibility with other pesticides can be tested by conducting a jar test with the volumes and rates typically used in agricultural application. Using a small container of water, add the proportionate amounts of the products: wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

The crop safety of all potential tank-mixes with VELUM RISE has not been tested on all crops listed on the label. Before applying any tank-mixture not specified on this label, safety to the target crop must be confirmed on a small portion of the crop listed on the label to be treated to ensure an adverse response will not occur.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

ROTATIONAL CROP RESTRICTIONS

The following crops can be replanted immediately following the last application of VELUM RISE:

Bulb vegetables (group 3-07); Cereals (groups 15 and 16) (except rice); Cotton (subgroup 20C); Dried shelled beans; Edible podded legume vegetables (subgroup 6A); Potato and other tuberous and corm vegetables (subgroup 1C); Rapeseed (subgroup 20A); Soybean; Succulent shelled pea and bean (subgroup 6B); Sugarbeet; Sunflower (subgroup 20B).

Alfalfa can be planted after 14 days.

The following crops can be planted after 30 days: Artichoke, (Globe); Brassica (cole) leafy vegetables (group 5); Citrus (group 10-10); Cucurbits (group 9); Dill seed; Fruiting Vegetables (group 8-10); Ginseng; Grapes and small vines (except fuzzy kiwifruit) (subgroup 13-07F); Herb (subgroup 19A); Hops; Leafy greens (subgroup 4A); Leaf petioles (subgroup 4B); Peanut; Pome fruit (group 11-10); Root vegetables (except sugar beet) (subgroup 1B); Small Berries, (canberries and bushberries) (subgroups 13-07A and 13-07B); Stone Fruits (group 12-12); Strawberry and other low-growing berries, except cranberry (subgroup 13-07G); Tobacco; Tree Nuts (group 14-12).

Sugarcane can be planted after 30 days in Region 3 (entire state of Florida; Mobile and Baldwin counties in Alabama).

DO NOT rotate to crops other than those listed above.

USE DIRECTIONS

POTATO		
Disease Suppression	Application Rate	Application Instructions
Stem & stolon canker and Tuber black scurf ¹ caused by soilborne <i>Rhizoctonia solani</i> .	13.0 fl oz/acre	Apply as an in-furrow spray at planting.
Black dot ¹ (<i>Colletotrichum coccodes</i>)	(0.212 lb/acre fluopyram)	In furrow spray - Mount spray nozzles to direct the spray as a 3- to 6-inch band down the row center onto the seed pieces in the furrow just before the seed is covered.
Early blight ¹ (<i>Alternaria solani</i>)	0.090 lb/acre penflufen)	
White mold ¹ (<i>Sclerotinia sclerotiorum</i>)		
Potato early dying/ Verticillium Wilt ¹ (<i>Verticillium dalia</i>)		
Pest Suppression	Application Rate	Application Instructions
Nematodes	13.0 fl oz/acre (0.212 lb/acre fluopyram) 0.090 lb/acre penflufen)	Apply as an in-furrow spray at planting. Alternatively, a single pre-plant incorporation application may be made. DO NOT make both a pre-plant incorporation and an in-furrow application with any fluopyram-containing products. In furrow spray - Mount spray nozzles to direct the spray as a 3- to 6-inch band down the row center onto the seed pieces in the furrow just before the seed is covered. Pre-plant incorporated – apply at a broadcast rate and incorporate. Despite suppression of root-knot nematode, tuber quality may not be adequately protected. If root-knot nematode is a severe economic pest, the use of other suppression measures should also be employed.

Restrictions:

- Maximum single application rate: 13.0 fl oz/acre VELUM RISE (0.212 lb/acre fluopyram and 0.090 lb/acre penflufen).
- Maximum annual application rate: 13.0 fl oz/acre VELUM RISE (0.212 lb/acre fluopyram and 0.090 lb/acre penflufen).
- Maximum number of applications per year: 1
- Regardless of formulation or method of application, **DO NOT** apply more than 0.446 lb fluopyram per acre per year (soil or foliar). **DO NOT** apply more than 0.22 lb fluopyram per acre per year before or at planting. **DO NOT** apply more than 0.143 lb penflufen per acre per year through combined seed-piece treatment and soil application.
- Apply using ground equipment.
- For soil application, to limit the potential for development of disease resistance to this chemical class, the first foliar fungicide application after VELUM RISE should be a product from a different FRAC group.
- Harvest at commercial maturity.

¹Not for use in California without a supplemental label.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross-contamination with other pesticides.

Pesticide Disposal: Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling:

Products in Non-Refillable Containers

Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Bayer, Bayer Cross and Velum are registered trademarks of Bayer Group.

©2025 Bayer Group. All rights reserved.

NET CONTENTS: 2 1/2 GALLONS



VELUM® RISE

FLUOPYRAM	GROUP	7	FUNGICIDE
PENFLUFEN	GROUP	7	FUNGICIDE

Fungicide and nematicide for use in potato as a soil treatment

ACTIVE INGREDIENTS:

Fluopyram*	22.12%
Penflufen*	9.38%
OTHER INGREDIENTS:	68.50%

TOTAL: 100.00%

Contains 2.08 lbs FLUOPYRAM and 0.88 lbs PENFLUFEN per U.S. gallon

*CAS Numbers 658066-35-4 (fluopyram) and 494793-67-8 (penflufen)

EPA Reg. No. 264-1222

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for First Aid Instructions and Booklet for Complete
Precautionary Statements and Directions for Use.

For MEDICAL and TRANSPORTATION Emergencies ONLY

Call 24 Hours a Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- **DO NOT** induce vomiting unless told to do so by a poison control center or doctor.
- Have person sip a glass of water if able to swallow.
- **DO NOT** give anything by mouth to an unconscious person.

IF ON SKIN or CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

In case of emergency, call the toll-free Bayer CropScience Emergency Response telephone number: 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Treat Symptomatically.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross-contamination with other pesticides.

Pesticide Disposal: Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Handling:

Products in Non-Refillable Containers

Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store, rinse for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture, and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Produced for
Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167

Bayer and Velum are registered trademarks of Bayer Group.
©2025 Bayer Group. All rights reserved.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE): Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, and waterproof or chemical-resistant gloves composed of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or viton ≥14 mils.

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- User should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

Bayer