

Wolverine[®] ADVANCED

Excellent Cross-Spectrum Weed Control in Barley

Wolverine[®] Advanced postemergent herbicide utilizes three different herbicide Groups for powerful, cross-spectrum weed control.

/// BENEFITS

- **Broad-Spectrum Weed Control**
Powered by herbicide Groups 27, 6 and 1 in an easy-to-use, all-in-one formulation.
- **Application Versatility**
Can be used on spring wheat, durum and barley with a wide application window.

/// WIDE APPLICATION WINDOW

- Best results are obtained when applications are made to young, actively growing weeds.
- Best applied on barley from emergence to 5-leaf stage.
- Best applied on wheat from emergence to 60 days prior to harvest in MN, MT, ND and SD, and 70 days prior to harvest in other states.
- Most effective on grass weeds from 1-leaf up to 2-tiller stage.
- Most effective on broadleaf weeds from one to six leaves (timing is species-dependent).

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.



LEARN MORE

Registered Crops

- Spring Wheat
- Durum
- Barley

Key Weeds Controlled

- Barnyard Grass
- Green Foxtail
- Annual Sowthistle
- Flix Weed
- Wild Buckwheat
- Wild Mustard
- Wild Oat
- Yellow Foxtail
- Cleavers
- Narrow-leaved Hawksbeard
- Perennial Sowthistle
- Lambsquarters
- Redroot Pigweed
- Russian Thistle
- Shepherdspurse
- Volunteer Canola
- Chickweed
- Common Ragweed
- Hemp-nettle
- Kochia

For a complete list of weeds controlled, visit CerealExperts.com

Application Rate

1.7 pt/A

Mode-of-Action Group

- 27 Pyrasulfotole
- 6 Bromoxynil
- 1 Fenoxoprop

Recommended Spray Volume

10 - 15 gal/A*

*For aerial application minimum spray volume of 5 gals/A.

Wolverine[®]
ADVANCED

*For more information on herbicide group numbers and mode-of-action options, visit IWillTakeAction.com